



ISSN 2278- 4136

ZDB-Number: 2668735-5

IC Journal No: 8192

Volume 1 Issue 6

Online Available at www.phytojournal.com



Journal of Pharmacognosy and Phytochemistry

Phytochemistry of Medicinal Plants

Mamta Saxena^{*1}, Jyoti Saxena¹, Rajeev Nema², Dharmendra Singh² and Abhishek Gupta²

1. Sarojini Naidu Government Girls Post Graduate (Autonomous) College, Shivaji Nagar, Bhopal - 462016 (M.P.), India. [E-mail: mamtasaxena00@yahoo.co.in]
2. Center for Microbiology & Bio-Technology Research and Training, Bhopal, India

Medicinal plants are a rich source of bioactive phytochemicals or bionutrients. Studies carried out during the past 2–3 decades have shown that these phytochemicals have an important role in preventing chronic diseases like cancer, diabetes and coronary heart disease. The major classes of phytochemicals with disease-preventing functions are dietary fibre, antioxidants, anticancer, detoxifying agents, immunity-potentiating agents and neuropharmacological agents. Each class of these functional agents consists of a wide range of chemicals with differing potency. Some of these phytochemicals have more than one function. There is, however, much scope for further systematic research in screening Indian medicinal plants for these phytochemicals and assessing their potential in protecting against different types of diseases

Keyword: Phytochemicals, Alkaloids, Terpenoids, Flavonoids, Saponins, Tannins and Phenolics.

1. Introduction

Phytochemicals (from the Greek word phyto, meaning plant) are biologically active, naturally occurring chemical compounds found in plants, which provide health benefits for humans further than those attributed to macronutrients and micronutrients^[1]. They protect plants from disease and damage and contribute to the plant's color, aroma and flavor. In general, the plant chemicals that protect plant cells from environmental hazards such as pollution, stress, drought, UV exposure and pathogenic attack are called as phytochemicals^[2,3]. Recently, it is clearly known that they have roles in the protection of human health, when their dietary intake is significant. More than 4,000 phytochemicals have been cataloged^[4] and are classified by protective function, physical characteristics and chemical characteristics^[5] and About 150 phytochemicals have been studied in detail^[4].

In wide-ranging dietary phytochemicals are found in fruits, vegetables, legumes, whole grains, nuts, seeds, fungi, herbs and spices^[3]. Broccoli, cabbage, carrots, onions, garlic, whole wheat bread, tomatoes, grapes, cherries, strawberries, raspberries, beans, legumes, and soy foods are common sources^[6]. Phytochemicals accumulate in different parts of the plants, such as in the roots, stems, leaves, flowers, fruits or seeds⁷. Many phytochemicals, particularly the pigment molecules, are often concentrated in the outer layers of the various plant tissues. Levels vary from plant to plant depending upon the variety, processing, cooking and growing conditions^[8]. Phytochemicals are also available in supplementary forms, but evidence is lacking that they provide the same health benefits as dietary phytochemicals^[4].

These compounds are known as secondary plant metabolites and have biological properties

such as antioxidant activity, antimicrobial effect, modulation of detoxification enzymes, stimulation of the immune system, decrease of platelet aggregation and modulation of hormone metabolism and anticancer property. There are more than thousand known and many unknown phytochemicals. It is well-known that plants produce these chemicals to protect themselves, but recent researches demonstrate that many phytochemicals can also protect human against diseases^[9].




Fig.1: Phytochemistry of medicinal plants.

Phytochemicals are not essential nutrients and are not required by the human body for sustaining life, but have important properties to prevent or to fight some common diseases. Many of these benefits suggest a possible role for phytochemicals in the prevention and treatment of disease. Because of this property; many researchers have been performed to reveal the beneficial health effects of phytochemicals. The purpose of the present review is to provide an overview of the extremely diverse phytochemicals presents in medicinal plants.

2. The Journey of Medicinal Plant Research

An assessment of the previous trends and impact of research into the phytochemistry on medicinal plants of the world is quite desirable before considering recent trends. After centuries of empirical use of herbal preparation, the first isolation of active principles alkaloids such as morphine, strychnine, quinine etc. in the early 19th century marked a new era in the use of medicinal plants and the beginning of modern medicinal plants research. Emphasis shifted away from plant derived drugs with the tremendous development of synthetic pharmaceutical chemistry and microbial fermentation after 1945. Plant metabolites were mainly investigated from a phytochemical and chemotaxonomic viewpoint during this period. Over the last decade, however, interest in drugs of plant and probably animal origin has grown steadily^[10]. Utilization of medicinal plants has almost doubled in Western Europe during that period. Ecological awareness, the efficacy of a good number of phytopharmaceutical preparations, such as ginkgo, garlic or valerian and increased interest of major pharmaceutical companies in higher medicinal plants as sources for new lead structures has been the main reasons for this renewal of interest. With the development of chemical science and pharmacognosy physicians began to extract chemical products from medicinal plants. A few examples of the products extracted from medicinal plants are - in 1920, quinine was isolated from Cinchona by the French pharmacist, Peletier & Caventou. In the mid-nineteenth century, a German chemist, Hoffmann obtained Aspirin from the bark of the willow. With the active principles in medicinal plants identified and isolated, plant-based prescriptions began to be substituted more and more with pure substances, which were more powerful and easier to prescribe and administer¹¹. Phytomedicine almost went into extinction during the first half of the 21st century due to the use of the 'more powerful and potent synthetic drug'. However, because of the numerous side effects of these drugs, the value of medicinal plants is being rediscovered as some of them have proved to be

as effective as synthetic medicines with fewer or no side effects and contraindications. It has been proved that although the effects of natural remedies may seem slower, the results are sometimes better on the long run especially in chronic diseases^[12].

3. Biological Activities of Phytochemicals

The phytochemicals present in plants are responsible for preventing disease and promoting health have been studied extensively to establish their efficacy and to understand the underlying mechanism of their action. Such studies have included identification and isolation of the chemical components, establishment of their biological potency both by *in vitro* and *in vivo* studies in experimental animals and through epidemiological and clinical-case control studies in man. Study findings suggest that phytochemicals may reduce the risk of coronary heart disease by preventing the oxidation of low-density lipoprotein (LDL) cholesterol, reducing the synthesis or absorption of cholesterol,

normalizing blood pressure and clotting, and improving arterial elasticity^[3,13]. Phytochemicals may detoxify substances that cause cancer. They appear to neutralize free radicals, inhibit enzymes that activate carcinogens, and activate enzymes that detoxify carcinogens. For example, according to data summarized by Meagher and Thomson, genistein prevents the formation of new capillaries that are needed for tumor growth and metastasis^[5]. The physiologic properties of relatively few phytochemicals are well understood and more many research has focused on their possible role in preventing or treating cancer and heart disease^[3]. Phytochemicals have also been promoted for the prevention and treatment of diabetes, high blood pressure, and macular degeneration^[4]. While phytochemicals are classified by function, an individual compound may have more than one biological function serving as both an antioxidant and antibacterial agent^[13]. Bioactive and Disease-preventing phytochemicals present in plant are shown in Table 1.

Table 1. Bioactive Phytochemicals In Medicinal Plants.

Classification	Main groups of compounds	Biological function
NSA (Non-starch poly-saccharides.)	Cellulose, hemicellulose, gums, mucilages, pectins, lignins	Water holding capacity, delay in nutrient absorption, binding toxins and bile acids
Antibacterial & Antifungal	Terpenoids, alkaloids, phenolics	Inhibitors of micro-organisms, reduce the risk of fungal infection
Antioxidants	Polyphenolic compounds, flavonoids, carotenoids, tocopherols, ascorbic acid	Oxygen free radical quenching, inhibition of lipid peroxidation
Anticancer	Carotenoids, polyphenols, curcumine, Flavonoids	Inhibitors of tumor, inhibited development of lung cancer, anti-metastatic activity
Detoxifying Agents	Reductive acids, tocopherols, phenols, indoles, aromatic isothiocyanates, coumarins, flavones, carotenoids, retinoids, cyanates, phytosterols	Inhibitors of procarcinogen activation, inducers of drug binding of carcinogens, inhibitors of tumourogenesis
Other	Alkaloids, terpenoids, volatile flavor compounds, biogenic amines	Neuropharmacological agents, anti-oxidants, cancer chemoprevention

4. Classification of Phytochemicals

The exact classification of phytochemicals could have not been performed so far, because of the wide variety of them. In recent year Phytochemicals are classified as primary or secondary constituents, depending on their role in plant metabolism. Primary constituents include the common sugars, amino acids, proteins, purines and pyrimidines of nucleic acids, chlorophyll's etc. Secondary constituents are the remaining plant chemicals such as alkaloids, terpenes, flavonoids, lignans, plant steroids, curcumines, saponins, phenolics, flavonoids and glucosides^[14]. Literature survey indicate that phenolics are the most numerous and structurally diverse plant phytoconstituents. (Figure 2).




Fig.2: Pie chart representing the major groups of plant Phytochemicals.

5. Phenolics

Phenolic phytochemicals are the largest category of phytochemicals and the most widely distributed in the plant kingdom. The three most important groups of dietary phenolics are flavonoids, phenolic acids, and polyphenols. Phenolic are hydroxyl group (-OH) containing

class of chemical compounds where the (-OH) bonded directly to an aromatic hydrocarbon group. Phenol (C₆H₅OH) is considered the simplest class of this group of natural compounds. Phenolic compounds are a large and complex group of chemical constituents found in plants^[15]. They are plant secondary metabolites, and they have an important role as defence compounds. phenolics exhibit several properties beneficial to humans and its antioxidant properties are important in determining their role as protecting agents against free radical-mediated disease processes. Flavonoids are the largest group of plant phenols and the most studied^[16]. Phenolic acids form a diverse group that includes the widely distributed hydroxybenzoic and hydroxycinnamic acids. Phenolic polymers, commonly known as tannins, are compounds of high molecular weight that are divided into two classes: hydrolyzable and condensed tannins.

6. Phenolic acids

The term “phenolic acids”, in general, designates phenols that possess one carboxylic acid functional group. Naturally occurring phenolic acids contain two distinctive carbon frameworks: the hydroxycinnamic and hydroxybenzoic structures (Figure3). Hydroxycinnamic acid compounds are produced as simple esters with glucose or hydroxy carboxylic acids. Plant phenolic compounds are different in molecular structure, and are characterized by hydroxylated aromatic rings^[17]. These compounds have been studied mainly for their properties against oxidative damage leading to various degenerative diseases, such as cardiovascular diseases, inflammation and cancer. Indeed, tumour cells, including leukaemia cells, typically have higher levels of reactive oxygen species (ROS) than normal cells so that they are particularly sensitive to oxidative stress^[18]. Many papers and reviews describe studies on bioavailability of phenolic acids, emphasizing both the direct intake through food consumption and the indirect bioavailability deriving by gastric, intestinal and hepatic metabolism^[19].

Table 2: The Major Classes of Phenolic Compounds in Plants

S.N.	Number of carbon atom	Basic skeleton	Class
1.	6	C ₆	Simple phenols Benzoquinones
2.	7	C ₆ -C ₁	Phenolic acids
3.	8	C ₆ -C ₂	Acetophenones Tyrosine derivatives
4.	9	C ₆ -C ₃	Hydroxycinnamic acid, Coumarins
5.	10	C ₆ -C ₄	Naphthoquinones
6.	13	C ₆ - C ₁ -C ₆	Xanthones
7.	14	C ₆ - C ₂ -C ₆	Stilbenes
8.	15	C ₆ - C ₃ -C ₆	Flavonoids
9.	18	(C ₆ - C ₃) ₂	Lignans
10.	30	(C ₆ - C ₃ -C ₆) ₂	Bioflavonoids
11.	N	(C ₆ - C ₃ -C ₆) _n	Condensed tannins

In addition Phenolic acid compounds and functions have been the subject of a great number of agricultural, biological, chemical and medical studies. In recent years, the importance of antioxidant activities of phenolic compounds and their potential usage in processed foods as a natural antioxidant compounds has reached a new level and some evidence suggests that the biological actions of these compounds are related to their antioxidant activity^[20].

5.1 Activity of Phenolic Acids

Phenolic compounds are famous group of secondary metabolites with wide pharmacological activities. Phenolic acid compounds and functions have been the subject of a great number of agricultural, biological, chemical and medical studies. Phenolic compounds in many plants are polymerized into larger molecules such as the proanthocyanidins (PA; condensed tannins) and lignins. Moreover, phenolic acids may arise in food plants as glycosides or esters with other

natural compounds such as sterols, alcohols, glucosides and hydroxyfatty acids. Varied biological activities of phenolic acids were reported. Increases bile secretion, reduces blood cholesterol and lipid levels and antimicrobial activity against some strains of bacteria such as staphylococcus aureus are some of biological activities of phenolic acids^[21]. Phenolics acid possesses diverse biological activities, for instance, antiulcer, anti-inflammatory, antioxidant^[22], cytotoxic and antitumor, antispasmodic, and antidepressant activities^[23].

6. Flavonoids

Flavonoids are polyphenolic compounds that are ubiquitous in nature. More than 4,000 flavonoids have been recognised, many of which occur in vegetables, fruits and beverages like tea, coffee and fruit drinks^[24]. The flavonoids appear to have played a major role in successful medical treatments of ancient times, and their use has persisted up to now. Flavonoids are ubiquitous

among vascular plants and occur as aglycones, glucosides and methylated derivatives. More than 4000 flavonoids have been described so far within the parts of plants normally consumed by humans and approximately 650 flavones and 1030 flavanols are known^[25]. Small amount of aglycones (i.e., flavonoids without attached sugar) are frequently present and occasionally represent a considerably important proportion of the total flavonoid compounds in the plant.

Figure 4, represents major flavonoids' structures. The six-membered ring condensed with the benzene ring is either -pyrone (flavones and

flavonols) or its dihydroderivative (flavanones and flavan-3-ols). The position of the benzenoid substituent divides the flavonoids into two classes: flavone (2-position) and isoflavone (3-position). Most flavonoids occur naturally associated with sugar in conjugated form and, within any one class, may be characterized as monoglycosidic, diglycosidic, etc. The glycosidic linkage is normally located at position 3 or 7 and the carbohydrate unit can be L-rhamnose, D-glucose, glucorhamnose, galactose or arabinose^[26].




Fig. 3. Structures of the important naturally occurring phenolic acids.

6.1 Activity of Flavonoids

Flavonoids have gained recent attention because of their broad biological and pharmacological activities in these order Flavonoids have been reported to exert multiple biological property including antimicrobial, cytotoxicity, anti-

inflammatory as well as antitumor activities but the best-described property of almost every group of flavonoids is their capacity to act as powerful antioxidants which can protect the human body from free radicals and reactive oxygen species. The capacity of flavonoids to act as antioxidants

depends upon their molecular structure. The position of hydroxyl groups and other features in the chemical structure of flavonoids are important for their antioxidant and free radical scavenging activities. On the other hand flavonoids such as luteolin and catechins, are better antioxidants than the nutrients antioxidants such as vitamin C, vitamin E and β -carotene. Flavonoids have been stated to possess many useful properties, containing anti-inflammatory

activity, enzyme inhibition, antimicrobial activity, oestrogenic activity, anti-allergic activity, antioxidant activity, vascular activity and cytotoxic antitumor activity^[27]. Flavonoids constitute a wide range of substances that play important role in protecting biological systems against the harmful effects of oxidative processes on macromolecules, such as carbohydrates, proteins, lipids and DNA^[28].




Fig.4. Chemical structures of some representative flavonoids.

7. Tannin

From a chemical point of view it is difficult to define tannins since the term encompasses some very diverse oligomers and polymers^[29,30]. It might be said that the tannins are a heterogeneous group of high molecular weight polyphenolic compounds with the capacity to form reversible and irreversible complexes with proteins (mainly), polysaccharides (cellulose, hemicellulose, pectin, etc.), alkaloids, nucleic acids and minerals, etc^[31,32,33]. On the basis of their structural characteristics it is therefore possible to divide the tannins into four major

groups: *Gallotannins*, *ellagitannins*, *complex tannins*, and *condensed tannins*^[34,35,36] (Figure 5).

(1) Gallotannins are all those tannins in which galloyl units or their *meta*-depsidic derivatives are bound to diverse polyol-, catechin-, or triterpenoid units.

(2) Ellagitannins are those tannins in which at least two galloyl units are C–C coupled to each other, and do not contain a glycosidically linked catechin unit.

(3) Complex tannins are tannins in which a catechin unit is bound glycosidically to a gallotannin or an ellagitannin unit.

(4) Condensed tannins are all oligomeric and polymeric proanthocyanidins formed by linkage of C-4 of one catechin with C-8 or C-6 of the next monomeric catechin.

Tannins are found commonly in fruits such as grapes, persimmon, blueberry, tea, chocolate,

legume forages, legume trees like *Acacia* spp., *Sesbania* spp., in grasses i.e; sorghum, corn, etc^[37]. Several health benefits have been recognized for the intake of tannins and some epidemiological associations with the decreased frequency of chronic diseases have been established^[38].




Fig.5. Classification of tannins.

7.1 Activity of Tannins

In medicine, especially in Asian (Japanese and Chinese) natural healing, the tannin-containing plant extracts are used as astringents, against diarrhoea, as diuretics, against stomach and duodenal tumours^[39], and as antiinflammatory, antiseptic, antioxidant and haemostatic pharmaceuticals^[40]. Tannins are used in the dyestuff industry as caustics for cationic dyes (tannin dyes), and also in the production of inks (iron gallate ink). In the food industry tannins are used to clarify wine, beer, and fruit juices. Other industrial uses of tannins include textile dyes, as antioxidants in the fruit juice, beer, and wine industries, and as coagulants in rubber Production⁴¹. Recently the tannins have attracted scientific interest, especially due to the increased

incidence of deadly illnesses such as AIDS and various cancers^{+42]}. The search for new lead compounds for the development of novel pharmaceuticals has become increasingly important, especially as the biological action of tannin-containing plant extracts has been well documented^[43,44].

8. Alkaloids

Alkaloids are natural product that contains heterocyclic nitrogen atoms, are basic in character. The name of alkaloids derives from the “alkaline” and it was used to describe any nitrogen-containing base^[45]. Alkaloids are naturally synthesis by a large numbers of organisms, including animals, plants, bacteria and fungi. Some of the fires natural products to be

isolated from medicinal plants were alkaloids when they first obtained from the plants materials in the early years of 19th century, it was found that they were nitrogen containing bases which formed salts with acid. Hence they were known as the vegetable alkalis or alkaloids and these alkaloids are used as the local anesthetic and stimulant as cocaine⁺^[46]. Almost all the alkaloids have a bitter taste. The alkaloid quinine for example is one of the bitterest tasting substances known and is significantly bitter (1×10^{-5}) at a molar concentration^[47].

Alkaloids are so numerous and involve such a variety of molecular structure that their rational classification is difficult. However, the best approach to the problem is to group them into families, depending on the type of heterocyclic ring system present in the molecule^[48]. For historical reasons as also because of their structural complexities, the nomenclature of alkaloids has not been systematized. The names of individual members are, therefore, generally derived from the name of the plant in which they occur, or from their characteristic physiological activity. The various classes of alkaloids according

to the heterocyclic ring system they contain are listed below.

Pyrrolidine alkaloids: they contain pyrrolidine (tetrahydropyrrole) ring system. E.g Hygrine found in *Erythroxylum coca* leaves.


Pyridine alkaloids: they have piperidine (hexahydropyridine) ring system. E.g Coniine, piperine and isopelletierine.

Pyrrolidine-pyridine alkaloids: the heterocyclic ring system present in these alkaloids is Pyrrolidine-pyridine. E.g Myosmine, Nicotine alkaloid found in tobacco (*Nicotiana tabacum*) plant.

Pyridine-piperidine alkaloids: This family of alkaloids contains a pyridine ring system joined to a piperidine ring system the simplest member is Anabasine alkaloid isolated from poisonous Asiatic plant anabasis aphyllan.

Quinoline Alkaloids: These have the basic heterocyclic ring system quinoline. E.g Quinine occurs in the bark of cinchona tree. It has been used for centuries for treatment of malaria. Synthetic drugs such as primaquine have largely replaced quinine as an anti-malarial.

Isoquinoline alkaloids: They contain heterocyclic ring system isoquinoline. E.g Opium alkaloids like narcotine, papaverine, morphine, codeine, and heroine.



Quinine [8]

Morphine [9]

Fig. 6. Structures of the important naturally occurring alkaloids.

8.1. Activity of Alkaloids

Alkaloids are significant for the protecting and survival of plant because they ensure their survival against micro-organisms (antibacterial and antifungal activities), insects and herbivores (feeding deterrents) and also against other plants by means of allelopathically active chemicals^[49]. The use of alkaloids containing plants as dyes, spices, drugs or poisons can be traced back almost to the beginning of civilization. Alkaloids have many pharmacological activities including antihypertensive effects (many indole alkaloids), antiarrhythmic effect (quinidine, sparteine), antimalarial activity (quinine), and anticancer actions (dimeric indoles, vincristine, vinblastine). These are just a few examples illustrating the great economic importance of this group of plant constituents^[50]. Some alkaloids have stimulant property as caffeine and nicotine, morphine are used as the analgesic and quinine as the antimalarial drug^[46].

9. Terpenoids

The terpenoids are a class of natural products which have been derived from five-carbon isoprene units. Most of the terpenoids have multi cyclic structures that differ from one another by their functional groups and basic carbon skeletons. These types of natural lipids can be found in every class of living things, and therefore considered as the largest group of natural products^[51]. Many of the terpenoids are commercially interesting because of their use as flavours and fragrances in foods and cosmetics examples menthol and sclareol or because they are important for the quality of agricultural products, such as the flavour of fruits and the fragrance of flowers like linalool^[52]. Terpenes

are widespread in nature, mainly in plants as constituents of essential oils. Their building block is the hydrocarbon isoprene, $\text{CH}_2=\text{C}(\text{CH}_3)-\text{CH}=\text{CH}_2$. Terpene hydrocarbons therefore have molecular formula $(\text{C}_5\text{H}_8)_n$ and they are classified according to the number of isoprene units^[53].

9.1 Hemiterpenoids: Consist of a single isoprene unit. The only hemiterpene is the Isoprene itself, but oxygen-containing derivatives of isoprene such as isovaleric acid and prenol is classified as hemiterpenoids.

9.2 Monoterpenoids: Biochemical modifications of monoterpenes such as oxidation or rearrangement produce the related monoterpenoids. Monoterpenoids have two isoprene units. Monoterpenes may be of two types i.e linear (acyclic) or contain rings e.g. Geranyl pyrophosphate, Eucalyptol, Limonene, Citral, Camphor and Pinene.

9.3 Sesquiterpenes: Sesquiterpenes have *three* isoprene units e.g. Artemisinin, Bisabolol and Farnesol, oil of flowers, or as cyclic compounds, such as Eudesmol, found in Eucalyptus oil.

9.4 Diterpenes: It composed for four isoprene units. They derive from geranylgeranyl pyrophosphate. There are some examples of diterpenes such as cembrene, kahweol, taxadiene and cafestol. Retinol, retinal, and phytol are the biologically important compounds while using diterpenes as the base.

9.5 Triterpenes: It consists of *six* isoprene units e.g. Lanosterol and squalene found in wheat germ, and olives.

9.6 Tetraterpenoids: It contains eight isoprene units which may be acyclic like lycopene, monocyclic like gamma-carotene, and bicyclic like alpha- and betacarotenes.




Fig. 7: Structures of the important terpenes of each class.

9.7 Activity of Terpenes

Among plant secondary metabolites terpenoids are a structurally most diverse group; they function as phytoalexins in plant direct defense, or as signals in indirect defense responses which involves herbivores and their natural enemies^[54]. Many plants produce volatile terpenes in order to attract specific insects for pollination or otherwise to expel certain animals using these plants as food. Less volatile but strongly bitter-tasting or toxic terpenes also protect some plants from being eaten by animals (antifeedants)^[55]. Last, but not least, terpenes play an important role as signal compounds and growth regulators (phytohormones) of plants, as shown by preliminary investigations. In addition, terpenoids can have medicinal properties such as anti-carcinogenic (e.g. perilla alcohol), antimalarial (e.g. artemisinin), anti-ulcer, hepatocidal, antimicrobial or diuretic (e.g. glycyrrhizin) activity and the sesquiterpenoid antimalarial drug artemisinin and the diterpenoid anticancer drug taxol.^[53,56]

10. Saponin

Saponins are a group of secondary metabolites found widely distributed in the plant kingdom. They form a stable foam in aqueous solutions such as soap, hence the name "saponin". Chemically, saponins as a group include compounds that are glycosylated steroids, triterpenoids, and steroid alkaloids. Two main types of steroid aglycones are known, spirostan and furostan derivatives (Figure 8A,B, respectively). The main triterpene aglycone is a derivative of oleanane (Figure 8C)^[57]. The carbohydrate part consists of one or more sugar moieties containing glucose, galactose, xylose, arabinose, rhamnose, or glucuronic acid glycosidically linked to a sapogenin (aglycone). Saponins that have one sugar molecule attached at the C-3 position are called monodesmoside saponins, and those that have a minimum of two sugars, one attached to the C-3 and one at C-22, are called bidesmoside saponins^[58].




Fig.8: Basic structure of steroid (A & B) and triterpenoid saponin (C)

10.1 Activity of Saponins

The physiological role of saponins in plants is not yet fully understood. While there is a number of a publication describing their identification in

plants, and their multiple effects in animal cells and on fungi and bacteria, only a few have addressed their function in plant cells. Many saponins are known to be antimicrobial, to inhibit

mould, and to protect plants from insect attack. Saponins may be considered a part of plants' defence systems, and as such have been included in a large group of protective molecules found in plants named phytoanticipins or phytoprotectants^[59]. Saponin mixtures present in plants and plant products possess diverse biological effects when present in the animal body. Extensive research has been carried out into the membrane-permeabilising, immunostimulant, hypocholesterolaemic and anticarcinogenic properties of saponins and they have also been found to significantly affect growth, feed intake and reproduction in animals. These structurally diverse compounds have also been observed to kill protozoans and molluscs, to be antioxidants, to impair the digestion of protein and the uptake of vitamins and minerals in the gut, to cause hypoglycaemia, and to act as antifungal and antiviral^[60,61,62].

11. Conclusion

Nature is a unique source of structures of high phytochemical diversity, many of them possessing interesting biological activities and medicinal properties. In the context of the worldwide spread different diseases such as AIDS, chronic diseases and a variety of cancers, an intensive search for new lead compounds for the development of novel pharmacological therapeutics is extremely important. With the present information are reported in this review, it is difficult to establish clear functionality and structure–activity relationships regarding the effects of phytochemicals in biological systems activity. This is largely due to the occurrence of a vast number of phytochemicals with similar chemical structures, and to the complexity of physiological reactions. Moreover, given the number of phytochemicals isolated so far, nature must still have many more in store. With the advances in synthetic methodology and the development of more sophisticated isolation and analytical techniques, many more of these phytochemicals should be identified.

12. Acknowledgement

The authors express gratitude Dr. Shobna Bajpai maroo, Principal Sarojini Naidu Government Girls (Post Graduate Autonomous) College and CMBT Laboratory, Bhopal, for kind support.

13. References

1. Hasler CM, Blumberg JB. Symposium on Phytochemicals: Biochemistry and Physiology. *Journal of Nutrition* 1999; 129: 756S-757S.
2. Gibson EL, Wardel J, Watts CJ. Fruit and Vegetable Consumption, Nutritional Knowledge and Beliefs in Mothers and Children. *Appetite* 1998; 31: 205-228.
3. Mathai K. Nutrition in the Adult Years. In Krause's Food, Nutrition, and Diet Therapy, 10th ed., ed. L.K. Mahan and S. Escott-Stump, 2000; 271: 274-275.
4. American Cancer Society. Phytochemicals. Available at http://www.cancer.org/eprise/main/docroot/ETO/content/ETO_5_3X_Phytochemicals, June 2000.
5. Meagher E, Thomson C. Vitamin and Mineral Therapy. In *Medical Nutrition and Disease*, 2nd ed., G Morrison and L Hark, Malden, Massachusetts: Blackwell Science Inc, 1999; 33-58.
6. Moorachian ME. Phytochemicals: Why and How? *Tastings*, 2000; 4-5.
7. Costa MA, Zia ZQ, Davin LB, Lewis NG. Chapter Four: Toward Engineering the Metabolic Pathways of Cancer-Preventing Lignans in Cereal Grains and Other Crops. In *Recent Advances in Phytochemistry*, vol. 33, Phytochemicals in Human Health Protection, Nutrition, and Plant Defense, ed. JT Romeo, New York, 1999; 67-87.
8. King A, Young G. Characteristics and Occurrence of Phenolic Phytochemicals. *Journal of the American Dietetic Association*, 1999; 24: 213-218.
9. Narasinga Rao. Bioactive phytochemicals in Indian foods and their potential in health promotion and disease prevention. *Asia Pacific Journal of Clinical Nutrition*, 2003; 12 (1): 9-22
10. Hamburger M, Hostettmann K. Bioactivity in Plants: The Link between Phytochemistry and Medicine. *Phytochemistry*, 1991; 30: 3864-3874.
11. Harvey A. Strategy for discovering drugs from previously unexploited natural products. *Drug Discovery Today*, 2000; 5: 294-300.
12. Akunyili DN. The role of regulation of medicinal plants and phytomedicine in socio-economic development, AGM/SC of the Nigerian Society of Pharmacognosy.
13. Abuja, 2003; 1-7.
14. Hahn NI. Is Phytoestrogens Nature's Cure for What Ails Us? A Look at the Research. *Journal of*

- the American Dietetic Association, 1998; 98: 974-976.
15. Walton NJ, Mayer MJ, Narbad A. Molecules of Interest: Vanillin. *Phytochemistry*, 2003; 63: 505-515.
 16. Dai J, Mumper R. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*, 2010; 15: 7313-7352.
 17. Balasundram N, Sundram K, Saman S. Phenolic Compounds in Plants and Agriindustrial by-Products: Antioxidant Activity, Occurrence, and Potential Uses. *Food Chemistry*, 2006; 99: 191-203.
 18. Mandal SM, Chakraborty D, Dey S. Phenolic acids act as signaling molecules in plant-microbe symbioses. *Plant Signal Behav*, 2010; 5: 359-368.
 19. Battisti V, Maders LD, Bagatini MD, Santos KF, Spanevello RM, Maldonado PA, Brulé AO, Araújo Mdo C, Schetinger MR, Morsch VM. Measurement of oxidative stress and antioxidant status in acute lymphoblastic leukemia patients. *Clin Biochem*, 2008; 41: 511-518.
 20. Lafay S and Gil-Izquierdo A. Bioavailability of phenolic acids. *Phytochemical Reviews*, 2008; 7: 301-311.
 21. Gryglewski RJ, Korbut R, Robak J. On the mechanism of antithrombotic action of flavonoids. *Biochemical Pharmacology*, 1987; 36: 317-321.
 22. Silva EM, Souza JNS, Rogez H, Rees JF, Larondelle Y. Antioxidant activities and polyphenolic contents of fifteen selected plant species from the Amazonian region. *Food Chemistry*, 2007; 101: 1012-18.
 23. Ghasemzadeh, A, Jaafar, HZE, Rahmat, A. Antioxidant activities, total Phenolics and flavonoids content in two varieties of Malaysia Young Ginger (*Zingiber officinale* Roscoe). *Molecules*, 2010; 15: 4324-4333.
 24. Pridham JB. In: *Phenolics in Plants in Health and Disease*, Pergamon Press, New York, 1960; 34-35.
 25. Harborne JB, Baxter H. *The handbook of natural flavonoids*, Volume 1 and 2. Chichester, UK: John Wiley and Sons, 1999.
 26. Pretorius JC. Flavonoids: A Review of Its Commercial Application Potential as Anti-Infective Agents. *Current Medicinal Chemistry-Anti Infective Agents*, 2003; 2: 335-353.
 27. Tapas AR, Sakarkar DM, Kakde RB. Flavonoids as Nutraceuticals: A Review. *Tropical Journal of Pharmaceutical Research*, 2008; 7: 1089-1099.
 28. Atmani D, Nassima C, Dina A, Meriem B, Nadjet D, Hania B, Flavonoids in Human Health: From Structure to Biological Activity. *Current Nutrition & Food Science*, 2009; 5: 225-237.
 29. Halliwell B, Gutteridge JMC. Role of free radicals and catalytic metal ions in human disease, An overview. *Methods Enzyme molecule*, 1990; 186: 1-85
 30. Harborne JB. An overview of antinutritional factors in higher plants. In: *Secondary plants products. Antinutritional and beneficial actions in animal feeding* Caygill JC and Mueller-Harvey I, eds. Nottingham Univ Press, UK, 1999; 7-16.
 31. Schofield P, Mbugua DM, Pell AN. Analysis of condensed tannins: a review. *Animal Feed Science Technology*, 2001; 91: 21-40.
 32. Vansoest PJ. *Nutritional ecology of the ruminant*, 2nd ed. Cornell Univ Press. Ithaca, NY, 1994; 476.
 33. Mueller-harvey I, Mcallan AB. Tannins: Their biochemistry and nutritional properties. *Advances in plant cell biochemistry and biotechnology*, Vol. 1 Morrison IM ed. JAI Press Ltd, London (UK), 1992; 151-217.
 34. Mangan JL. Nutritional effects of tannins in animal feeds. *Nutrition Research and Reviews*, 1988; 1: 209-231.
 35. Mc-Leod MN. Plant tannins: Their role in forage quality. *Nutrition Abstract Review*, 1974; 44: 803-812.
 36. Mole S, Waterman PG. Tannic acid and proteolytic enzymes: enzyme inhibition or substrate deprivation? *Phytochemistry*, 1987; 26: 99-102.
 37. Giner-Chavez BI. *Condensed tannins in tropical forages*. Doctoral Thesis. Cornell University. Ithaca, NY, 1996.
 38. Serrano J, Puupponen-Pimia R, Dauer A, Aura A, Saura-Calixto F. Tannins: current knowledge of food sources, intake, bioavailability and biological effects. *Molecular Nutrition Food Research*, 2009; 53: S310-29.
 39. De Bruyne T, Pieters L, Deelstra H, Vlietinck A. Condensed vegetables tannins: biodiversity in structure and biological activities. *Biochemical System Ecology*, 1999; 27: 445-59.
 40. Dolara P, Luceri C, De Filippo C, Femia AP, Giovannelli L, Cardeni G, Cecchini C, Silvi S, Orpianesi C, Cresci A. Red wine polyphenols influence carcinogenesis, intestinal microflora, oxidative damage and gene expression profiles of colonic mucosa in F344 rats. *Mutation Research*, 2005; 591: 237-46.
 41. Gyamfi MA, Aniya Y. Antioxidant properties of Thonningianin A, isolated from the African medicinal herb, *Thonningia sanguine*. *Biochemical Pharmacology*, 2002; 63: 1725-37.
 42. Blytt HJ, Guscar TK, Butler LG. Antinutritional effects and ecological significance of dietary condensed tannins may not be due to binding and inhibiting digestive enzymes. *Journal of Chemical Ecology*, 1988; 14: 1455-1465.
 43. Palavy K, Priscilla MD. Standardisation of selected Indian medicinal herbal raw material

- containing polyphenols as major constituents. *Journal Pharmaceutical sciences*, 2006; 68: 506-509.
44. Mueller-Harvey I. Tannins: their nature and biological significance. In: *Secondary plants products. Antinutritional and beneficial actions in animal feeding* Caygill JC and Mueller-Harvey I, eds. Nottingham Univ Press (UK), 1999; 17-70.
 45. Mueller-Harvey I, McAllan AB. Tannins. Their biochemistry and nutritional properties. In: *Advances in plant cell biochemistry and biotechnology*, Vol. 1 Morrison IM, ed. JAI Press Ltd, London (UK), 1992; 151-217.
 46. Rao RVK, Ali N, Reddy MN. Occurrence of both saponin and alkaloid lycorine in *Curculigo orchioides*. *Indian Journal Pharma Science*, 1978; 40: 104-105.
 47. Mishra SN. Analytical methods for analysis of total alkaloids in root of *Withania* spp. *Proc. All India workshop on M&AP*, Faizabad, 1989; 492-95.
 48. Krishnan R, Chandravada MV, Ramachander PR, Bharathkumar H. Inter-relationships between growth and alkaloid production in *Catharanthus roseus* G. Don. *Herba Hungarica*, 1983; 22: 47-54.
 49. Molyneux RJ, Nash RJ, Asano N. Alkaloids: Chemical and Biological Perspectives, Vol. 11, Pelletier SW, ed. Pergamon, Oxford, 1996; 303.
 50. Wink M, Schmeller T, Latz-Briining B. Modes of action of allelochemical alkaloids: Intraction with neuroreceptors, DNA and other molecular targets. *Journal of chemical Ecology*, 1998; 24: 1888-1937.
 51. Elbein AD, Molyneux RJ. *Comprehensive Natural Products Chemistry*, Vol. 3, Barton D and Nakanishi K, ed. Amsterdam, 1999; 129.
 52. Harborne JB, Tomas-Barberan FA. *Ecological Chemistry and Biochemistry of Plant Terpenoids*, Clarendon, Oxford, 1991.
 53. Langenheim JH. Higher plant terpenoids: A phytocentric overview of their ecological roles. *Journal of Chemical Ecology*, 1994; 20: 1223-1280.
 54. McCaskill D, Croteau R. Some caveats for bioengineering terpenoid metabolism in plants. *Trends Biotechnology*, 1998; 16: 349-355
 55. Degenhardt J, Gershenzon J, Baldwin IT, Kessler A. Attracting friends to feast on foes: Engineering terpene emission to make crop plants more attractive to herbivore enemies. *Current Opinion Biotechnology*, 2003; 14: 169-176.
 56. Dudareva N, Pichersky E, Gershenzon J. Biochemistry of plant volatiles. *Plant Physiology*, 2004; 135: 1893-1902.
 57. Bohlmann J, Meyer-Gauen G, Croteau R. Plant terpenoid synthases: Molecular biology and phylogenetic analysis. *Proc Natl Acad Sci USA*, 1998; 95: 4126-4133.
 58. Lasztity R, Hidvegi M, Bata A. Saponins in food. *Food Review International*, 1998; 14: 371-390.
 59. Lacaille-Dubois MA, Wagner H. Bioactive saponins from plants: An update. In *Studies in Natural Products Chemistry*; Atta-Ur-Rahman, ed. Elsevier Science. Amsterdam, 2000; 21: 633-687.
 60. Morrissey JP, Osbourn AE. Fungal resistance to plant antibiotics as a mechanism of pathogenesis. *Microbiological and Molecular Biological Reviews*, 1999; 63: 708-724.
 61. Takechi M, Matsunami S, Nishizawa J, Uno C, Tanaka Y. Haemolytic and antifungal activities of saponins or anti-ATPase and antiviral activities of cardiac glycosides. *Planta Medica*, 1999; 65: 585-586.
 62. Traore F, Faure R, Ollivier E, Gasquet M, Azas N, Debrauwer L, Keita A, Timon-David P, Balansard G. Structure and antiprotozoal activity of triterpenoid saponins from *Glinus oppositifolius*. *Planta Medica*, 2000; 66: 368-371.
 63. George F, Zohar K, Harinder PS, Makkar, Klaus B. The biological action of saponins in animal systems: a review. *British Journal of Nutrition*, 2002; 88: 587-605.