



ISSN 2278-4136
ISSN 2349-8234
JPP 2014; 2 (6): 53-56
Received: 01-02-2014
Accepted: 12-01-2014

Dr. Mahesh Chand Meena,
Lecturer in Botany,
Rajesh Pilot Govt. PG College,
Lalsot (Dausa) Rajasthan,
Tel: +91-9413090517
E-mail: meenamc@rediffmail.com

Dr. Vidya Patni,
Associate Professor,
University of Rajasthan,
Jaipur (Raj.)
E-mail: patni-vidya@uniraj.ernet.in
Tel: +91-9829059898

Dr. Rishi Kesh Meena
Assistant Professor,
University of Rajasthan,
Jaipur (Raj.)
E-mail: rishi_1180@yahoo.com
Tel: +91-9828175740

Correspondence:

Dr. Mahesh Chand Meena,
Lecturer in Botany,
Rajesh Pilot Govt. PG College,
Lalsot (Dausa) Rajasthan,
Tel: +91-9413090517
E-mail: meenamc@rediffmail.com

High frequency plant regeneration from shoot tip explants of *Citrullus colocynthis* (Linn.) Schrad. – An important medicinal herb

Mahesh Chand Meena, Rishi Kesh Meena and Vidya Patni

ABSTRACT

A high frequency and rapid regeneration protocol was developed from shoot tip explants of *Citrullus colocynthis* on MS medium supplemented with BAP (0.5 mg/l) and NAA (0.5 mg/l). Highest number of shoots (23.0 ± 0.567) was obtained on MS medium containing BAP (0.5 mg/l) and NAA (0.5 mg/l). The regenerated shoots were further elongated on same medium. *In vitro* shoots were excised from shoot clumps and transferred to rooting medium containing IBA (4.0 mg/l) with 0.2% activated charcoal. The rooted plants were hardened in polycups containing sterile soil and vermiculate and finally well established in the field; survival rate was 60%. This is the first report of direct *in vitro* plantlet regeneration in *Citrullus colocynthis* from shoot tip explant.

Keywords: *Citrullus colocynthis*, Medicinal Plant, Regeneration, Shoot tip.

1. Introduction

Plant based remedies have always been an integral part of traditional medicine throughout the world. The demand for herbal remedies has been increasing significantly^[1]. Constant demands for some of the medicinal plants have affected their availability to a large extent. Tissue culture techniques are being increasingly exploited for clonal multiplication and *in vitro* conservation of valuable indigenous germplasm threatened with extinction. Greater demand for these plants especially for the purpose of food and medicine is one of the causes of their rapid depletion from primary habitats. Micropropagation offers a great potential for large scale multiplication of such useful species and subsequent exploitation^[2].

Among the important medicinal plants, *Citrullus colocynthis* (Linn.) Schrad (Cucurbitaceae) commonly known as, Tumba or Indrayan or Bitter apple, is a perennial herb and a common desert plant having a wide range growing in the most barren and arid situations. *C. colocynthis* has been utilized as a general promoter of health in medicine.

C. colocynthis is used in folk medicine by people in rural areas as purgative, antirheumatic, anthelmintic, anticancerous and as remedy for skin infection. Pulp of fruit has antibacterial activity. The dried and powdered pulp is taken orally to cause abortion. Seeds are purgative and leaves are used against migraine and neuralgia. Root has a beneficial action in ophthalmia, uterine pains, jaundice and is also useful against boils, pimples and enlarged abdomen.

Due to unsuitable exploitation of eco-resources, several plant and tree species of great medicinal importance have become threatened and may become extinct due to lack of proper planning for their conservation and multiplication. Due to excessive and destructive exploitation of *C. colocynthis*, it is getting fast depleted^[3]. Therefore, it is time to conserve this plant species by means of micropropagation through tissue culture technique. Tissue culture is an age-old practice for *in vitro* regeneration of plants, especially the economically valuable plant or plants that are difficult to propagate in natural environment. This method is also used for studying the plant development at the molecular level thereby artificially increasing the plant output molecules. So we describe here an efficient procedure for the rapid clonal propagation of *C. colocynthis* through shoot tip culture, which is often used to produce disease free stock of plants. Shoot tip culture is widely used for rapid propagation of many species due to its advantages over traditional methods^[4]. The main objective of the present investigation was to select the best *in vitro* responding media combinations for inducing morphogenesis in higher frequencies and optimization of conditions for complete plantlet development.

2. Materials and Methods

Shoot tip bearing shoot segments were collected from mature plants growing in University campus, Jaipur. Explants were initially washed under running water with Labolene solution (0.1% v/v) for 5 minutes and sterilized with 0.05% (w/v) mercuric chloride for 2-3 min. Shoot tips were rinsed with sterile distilled water 4-5 times to remove the traces of mercuric chloride. These shoot tips were inoculated on MS-medium supplemented with different concentrations and combinations of phytohormones. All the media contained 3% (w/v) sucrose. The pH of the media was adjusted to 5.8 and solidified with 0.8% agar, before autoclaving at a pressure of 15 psi and 121 °C for 20 min. All the cultures were maintained at a temperature of 25 ± 2 °C under a 16/8-h (light/dark) photoperiod provided by cool white fluorescent light. Multiple shoots obtained were later transferred to elongation medium supplemented with different growth hormones. The cultures were

regularly subcultured on fresh medium at four weeks intervals and observations were recorded. Elongated and healthy plantlets were transferred to rooting medium containing (4.0 mg/l) concentration of IBA with activated charcoal (0.2%). Regenerated plantlets were then transferred to pots containing sterilized soil and vermiculite (3:1). The plantlets were kept in a glass house for one week under normal light and temperature. New leaves emerged from the plantlets after 10-15 days in the glass house; they were then placed in the normal environment for 1h and assessed for signs of wilting. The exposure time was increased daily until the plants were established fully under normal environment conditions, they were then transferred into the field for their normal growth. For each treatment 8 replicates were used and each experiment was repeated at least five times. The cultures were examined periodically. The means and standard errors of the results were also calculated.

Table 1: Effect of Plant Growth Regulators (PGR) on shoot formation from shoot tip explant of *Citrullus colocynthis* on MS-medium

S. No.	PGR's mg/l			No. of shoots buds per explant Mean ±SE
	BAP	Kn	NAA	
1.	0.5	-	-	5.6 ± 0.220
2.	1.0	-	-	7.0 ± 0.756
3.	2.0	-	-	11.2 ± 0.180
4.	3.0	-	-	10.2 ± 0.336
5.	4.0	-	-	8.6 ± 0.220
6.	5.0	-	-	6.5 ± 0.220
7.	-	0.5	-	2.4 ± 0.220
8.	-	1.0	-	3.6 ± 0.359
9.	-	2.0	-	5.8 ± 0.595
10.	-	3.0	-	8.2 ± 0.336
11.	-	4.0	-	6.6 ± 0.608
12.	-	5.0	-	4.2 ± 0.336
13.	0.5	0.5	-	7.0 ± 0.401
14.	1.0	1.0	-	9.4 ± 0.220
15.	1.5	1.5	-	13.4 ± 0.463
16.	2.0	2.0	-	11.0 ± 0.284
17.	2.5	2.5	-	7.4 ± 0.588
18.	3.0	3.0	-	7.2 ± 0.336
19.	0.5	-	0.5	23.0 ± 0.567
20.	1.0	-	1.0	16.8 ± 0.180
21.	1.5	-	1.5	14.6 ± 0.220
22.	2.0	-	2.0	12.8 ± 0.336
23.	2.5	-	2.5	10.0 ± 0.634
24.	3.0	-	3.0	7.8 ± 0.595

Table 2: Effect of auxin (IBA) along with 0.2% activated charcoal on root induction from shoots of *C. colocynthis* after 4 weeks of culture. [values are mean ± SE from 5 replicates in each treatment]

IBA (mg/l)	% of cutting rooted	No. of roots per shoots	Average length of the root (cm)
1.0	80	1.8 ± 0.456	2.0 ± 0.16
2.0	70	8.0 ± 0.283	2.2 ± 0.18
3.0	75	13.8 ± 0.335	2.5 ± 0.40
4.0	95	24.0 ± 1.678	3.0 ± 0.16
5.0	85	16.0 ± 0.671	2.6 ± 0.63

3. Results and Discussion

Shoot tip of Indrayan (*Citrullus colocynthis*) were isolated

aseptically and cultured on MS-medium supplemented with cytokinins and auxins for initiating vegetative growth and inducing

maximum number of plantlets.



Fig 1: *In vitro* shoot multiplication and acclimatization of *Citrullus colocynthis* through shoot tip culture. A. Shoot tip explant on MS medium. B&C. Multiple shoot induction after two weeks and three weeks on MS medium supplemented with BAP (0.5 mg/l) and NAA (0.5 mg/l). D. Elongated shoots on MS medium supplemented with BAP (0.5 mg/l) and NAA (0.5 mg/l). E. *In vitro* rooting of regenerated shoot on MS medium supplemented with IBA (4.0 mg/l) and 0.2 % activated charcoal. F. Hardening of *in vitro* raised plantlets.

The shoot tip explants were inoculated on MS medium containing different concentrations of BAP (0.5-5.0 mg/l) and Kn (0.5-5.0 mg/l) alone and in combination (0.5-3.0 mg/l BAP and 0.5-3.0 mg/l Kn and 0.5-3.0 mg/l BAP and 0.5-3.0 mg/l NAA) (Fig. 1A). Among different concentrations used best response towards shoot proliferation from shoot tip explant was obtained on MS medium with BAP (0.5 mg/l) and NAA (0.5 mg/l). Initiation of axillary bud started after one week of inoculation (Fig. 1B). Maximum multiple shoots (23.0 ± 0.567) were obtained on MS medium supplemented with BAP (0.5 mg/l) and NAA (0.5 mg/l) (Fig. 1C). Shoots elongated on the same medium (Fig. 1D). From the results (Table 1) it is clear that a combination of BAP (0.5 mg/l) and NAA (0.5 mg/l) at lower concentration was suitable for shoot multiplication as well as shoot elongation.

The effective role of NAA in combination with BAP for the induction of multiple shoots has been reported in many plant species [5-8]. In contrast to above mentioned results some researchers observed that the combination of BAP and IAA on MS-medium favoured multiple shoot buds in some plant species [9-10]. Combination of cytokinins also favoured multiple shoot proliferation in many plant species [11-13]. Rooting of the micropropagated plants was usually achieved in auxin containing medium [14]. Root formation is an energy demanding process and thus exogenous supply of carbohydrates is required. However this being the last stage of *in vitro* culture it is important to transform the plant from heterotrophic to autotrophic mode of nutrition. Thus the supply of exogenous sugars should be

reduced at this time. The rooting response differed according to different concentrations and combinations of auxins used (Table.2). In the present study, rooting was achieved on full strength of MS-medium fortified with IBA (4.0 mg/l) and activated charcoal (0.2 %). On this medium, rooting percentage (95 %), number of roots (24.0 ± 1.678) and root length (3.0 ± 0.16) were maximum. Roots were thick and with white root hairs. Similar effect of IBA on rooting in several medicinal plant species were also observed by several workers in different plant species [15-20].

Plantlets with 6-7 leaves and well developed root system were removed and transferred to pot containing soilrite (Fig. 1F). These pots were kept in growth chamber for 15 days at 26 ± 2 °C and 2000 lux intensity for acclimatization. In order to maintain high humidity the pots were covered with inverted glass beaker. After six months when new leaves emerged from these plantlets, they were taken outside the growth chambers and kept in shady place under normal temperature and light. A 60 % survival rate was obtained when acclimatized plantlets were transferred to green house.

4. Conclusion

The present paper describes a prime and easy-to-use protocol for large scale production of plantlets through shoot tip culture of *Citrullus colocynthis* and the method is useful for the *ex-situ* conservation of other threatened medicinally important species. This is the first report of *in vitro* culture of this wild endangered medicinal herb. So the result of present investigation forms the basis of further research. Since, this plant has soil binding property and multiplied in bulk, it can be used to control soil erosion as well as spreading of Thar Desert in Rajasthan. Many wild cultivated species are the major sources of genes for making hybrid varieties (Turtumba with *Citrullus vulgaris* and *C. colocynthis*). *Citrullus colocynthis* being an endangered wild medicinal herb has to be propagated and conserved. This work is a part of the efforts taken towards that direction.

5. Acknowledgement

The senior author (Mahesh Chand Meena) wishes to acknowledge University Grant Commission, CRO-Bhopal for financial support.

6. References

- Suchitra B, Mehar Z, Susil K. *In vitro* multiplication of *Centella asiatica*. Current Science 1999; 76:147-158.
- Boro PS, Shirma Deka AC, Kalita MC. Clonal propagation of *Alternanthera sessilis*: A biopharmaceutically potent herbal medicinal plant. J. Phytol. Res. 1998; 11:103-106.
- Bhandari, MM. Medicinal plants: Biodiversity conservation, export potential and intellectual property rights. In: Medicinal plants utilization and conservation (Eds.) Trivedi P.C., Aviskar Publishers, Jaipur. 2004; 83.
- Hu CY, Wang PJ. Meristem, shoot tip and bud cultures. In: Handbook of Plant Cell Culture: Techniques for Propagation and Breeding. Vol. I (Eds.) Evans, D.A., Sharp, W.R., Ammirato, P.V. and Yamada, Y. Macmillan Publishing Co., New York, 1983; 177-227.
- Chakraborty D, Bandhyopadhyay A, Bhattacharyya PK, Gupta K. *In vitro* flowering of *Basilicum polystachyon* (L.) Moench. *Plant Cell Biotech. and Mol. Biol.* 2006; 7(1&2):17-22.
- Kalimuthu K, Saravankumar M, Senthilkumar R. *In vitro* micropropagation of *Musa sapientum* L. (Cavendish Dwarf). *African Journal of Biotechnology.* 2007; 6(9):1106-1109.
- Baksha R, Jahan MAA, Khatun R, Munshi, JL. *In vitro* rapid clonal propagation of *Rauwolfia serpentina* (Linn.). *Bangladesh J. Sci. Ind. Res.* 2007; 42(1):37-44.
- Karuppusamy S, Pullaiah T. *In vitro* shoot multiplication of *Bupleurum distichophyllum* Wight – A native medicinal plant of Southern India. *Plant Tissue Cult. & Biotech.* 2007; 17(2):115-124.
- Sobhakumari VP, Lalithakumari D. Direct plant regeneration from shoot tip cultures of *Capsicum annuum* L. cv. PLR-1. *Phytomorphology.* 2003; 53(3&4): 235-242.
- Sharma MM, Jalootharia D, Khanna P, Batra A. An efficient *in vitro* mass propagation of a medicinally potent plant species *Vitex negundo* L. viz nodal segments. *Phytomorphology.* 2006; 56(1&2): 35-39.
- Soni V. Morphogenetic, biochemical and physiological studies in *Commiphora wightii*, Ph.D. Thesis, University of Rajasthan, Jaipur. 2005
- Girija S, Kavitha S, Deepavathi S. Direct multiple shoot regeneration from shoot tip and nodal explants of *Ocimum sanctum* L. (Tulsi): A medicinal herb. *Plant Cell Biotech. Mol. Biol.* 2006; 7(1&2): 23-28.
- Rao S, Pratibha GS, Parashuram YJ, Kaviraj CP. High frequency plant regeneration from shoot tip explants of chilli (*Capsicum annuum*). *Plant cell Biotechnol. Mol. Biol.* 2006; 7:163-166.
- Gaspar T, Coumans M. Root formation. In: Cell and tissue culture in forestry. Vol.II Bong, J.M. and Durzan, D.J. (Eds.). Marlinus Nijhof Publishers Dordrecht. 1987; 202-217.
- Sunandakumari C, MaSurtin KP, Chithra M, Sini S, Madhusoodanan PV. Rapid axillary bud proliferation and *ex vitro* rooting of herbal spice *Mentha piperita* L. *Indian J Biotechnol.* 2004; 3: 108-112.
- Sivanesan I, Murugesan K. *In vitro* adventitious shoot formation from leaf explants of *Withania somnifera* Dunal. *Plant Cell Biotechnol. Mol. Biol.* 2005; 6: 163-166.
- Chakraborty S, Roy SC. Micropropagation of *Cyphomandra betacea* (Cav.) Sendt., A potential horticultural and medicinal plant, by axillary bud multiplication. *Phytomorphology* 2006; 56(1&2):29-33.
- Sharma A, Meena MC, Patni V, Arora DK (2007) *In vitro* multiplication of *Acalypha wilkesiana* 'dwarf' through nodal segment culture. *Jour. Pl. Sci. Res.* 2007; 23 (1-2):79-82.
- Gadidasu K, Umate P, Aileni M, Kota SR, Kokkiralala VR, Kasula K. Micropropagation of a valuable ethnomedicinal plant *Streblus asper* Lour. *J Phytol* 2011; 3(2):18-23.
- Hossain MJ, Khaleda L, Al-Forkan M. Development of an efficient *In vitro* micropropagation protocol for medicinally important plant *Achyranthes bidentata* Blume. *Journal of Pharmacognosy and Phytochemistry* 2013; 2(4):6-13.