



E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2019; 8(1): 764-769
Received: 02-11-2018
Accepted: 06-12-2018

Sanjay Sapkal
Division of Agricultural
Economics, IARI, New Delhi,
India

BH Kamble
Department of Agricultural
Economics, MPKV, Rahuri,
Maharashtra, India

Pramod Kumar
Division of Agricultural
Economics, IARI, New Delhi,
India

Amit Kar
Division of Agricultural
Economics, IARI, New Delhi,
India

GK Jha
Division of Agricultural
Economics, IARI, New Delhi,
India

Correspondence
Sanjay Sapkal
Division of Agricultural
Economics, IARI, New Delhi,
India, India

Impact of laser land levelling in rice-wheat systems of the North-Eastern indo-gangetic plains of India

Sanjay Sapkal, BH Kamble, Pramod Kumar, Amit Kar and GK Jha

Abstract

The paper has compared the economics of wheat as well as paddy production in North-Eastern Indo-Gangetic Plains of Bihar under laser land levelling and conventional methods and assessed the contribution of technology and inputs to the increased productivity due to laser land levelling (LLL). The net income has been found higher in LLL method, mainly due to incremental gain in productivity compared to that in conventional method. The study has observed that LLL technology has potential to provide additional income to farmers and help in conservation of scarce resources. The decomposition of productivity revealed 17 per cent and 47 per cent of the difference in productivity of wheat and paddy, respectively due to LLL and the rest due to changes in input use. In all, about Rs.20055 of net incremental benefit was derived from per LLL operation. Despite several economic and environmental advantages, adoption of LLL technology has been limited and one major constraint identified as the small size of land holding followed by lack of technical knowledge. The study has suggested that LLL technology should be disseminated on a wider scale with the help of co-operative farming and better technical as well as extension support to the farmers.

Keywords: Laser land levelling, conventional tillage, economics, decomposition analysis

Introduction

With increasing climate variability and rapid melting of glaciers, water scarcity is expected to be a major challenge to agricultural production and food security in South Asia. Increasing temperature, which has been observed in most of the region, increases the demand for irrigation water. The groundwater table is observed to be declining in India (Aggarwal *et al.* 2004; Joshi and Tyagi 1994; Kerr 2009; Kumar *et al.* 2007) [2, 17, 19, 20]. As agriculture is the largest beneficiary of groundwater, a declining groundwater table will impact the sustainability of agriculture and the overall food security of the country. Furthermore, the use of motorised pump to extract groundwater for irrigation raised the demand for energy by the agriculture sector in India. Given the existing consumption pattern, India will need to produce at least 37 per cent more rice and wheat by 2025 as compared to that of the year 2000, with nearly 10 per cent less water that will be available for irrigation (Jat *et al.* 2006) [14]. Therefore, there is a need of technologies that reduces water requirement, increasing the efficiency of energy use and enhances agricultural productivity. But, in the rice-wheat (RW) system of the Indo-Gangetic Plains (IGP), about 10–25 per cent of irrigation water is lost due to poor water management and uneven fields (Kahlowan *et al.* 2000) [18]. To overcome such a situation, the laser land levelling (LLL) is a technology that has the primary benefit of reduction in the loss of irrigation water occurring due to highly undulating land with achieving the desired level of accuracy through laser equipped drag buckets. It also facilitates uniformity in the placement of seeds or seedlings and promotes better crop stands, which eventually contributes to higher crop yields. A uniform field improves irrigation efficiency through better control of water distribution and reduces the potential for nutrient loss through improved runoff control, leading to greater efficiency of fertilizer use and higher yields. Despite this, there has been no comprehensive study which examines the impact of LLL on crop yield, irrigation duration, energy saving, and overall farm profitability. Hence, the major focus of the study is to assess the impact of LLL on cost of inputs, crop yield, irrigation duration, energy saving, and farm profitability.

Methodology

Laser land levelling (LLL) has been interpreted here as the process of levelling the land with the help of laser land leveller. The conventional tillage (CT) refers to the intensive tillage with multiple passes of a tractor to accomplish land preparation. LLL technology was found to be more skewed towards medium and large farmers in Bihar.

For this study, Vaishali district was selected purposively due to implementation of climate smart village project by CGIAR through CIMMYT and other co-partners. From the Vaishali district, four climate smart villages (CSV) and four non-climate smart villages (Non-CSV) were selected. A total of 30 farmers from CSV who adopted laser land levelling technology for wheat and paddy production were selected randomly. From the Non-CSV villages, 40 numbers of farmers practising conventional tillage method were selected. The characteristics and socio-economic conditions of both types of the households were almost similar. The primary data were collected during the agriculture year 2016-17 from 70 farmers.

The farm management cost concept was used to analyse the impact of LLL technology on farm income. The sample farmers were classified into two category based on their adoption of technology i.e. LLL technology adopter and non-adopter. The impact of the adapted technology was assessed in terms of enhancement in yield, income, and improvement in efficiency. In which the various cost components such as Costs A1, B1, B2, C2, and C3 was calculated to analyse the profitability of the wheat and paddy production.

Cost A1= Wages of hired labour, cost of inputs such as seed, manures, liquid and solid fertilizers, insecticides and pesticides, hired machinery charges, imputed value of owned machine power, implements like fertigation units and farm buildings, irrigation charges, land revenue and interest on working capital.

Cost B1 = Cost A1 + Interest on the value of owned fixed capital assets (excluding land).

Cost B2 = Cost B1 + Rental value of land.

Cost C1 = Cost B1 + Imputed value of family labour.

Cost C2 = Cost B2 + Imputed value of family labour.

Cost C3 = Cost C2 + 10 percent of Cost C2 accounting for managerial input.

The cost C_1 was taken into account in this study to calculate net income and benefit-cost ratio. The cost of irrigation was calculated by multiplying the time required to irrigate the farm with cost of electricity or diesel consumption per hour. The cost of electricity was taken based on per unit rate fixed by the Bihar Electricity Distribution Corporation. The cost on human labour, machine labour and diesel were taken on actual expenditure basis. Gross income included the total value of main crop and by-products. Net income was calculated as the difference between gross income and cost of production (cost C_1).

The output decomposition model developed by Bisaliah (1977) [8] was used to quantify the contribution of various sources to the productivity differences between LLL and CT methods. It is expected that the practice of LLL technology will result in changes in input-use pattern, which in turn will affect the land productivity. Hence, increase in land productivity is not only due to adoption of LLL method but also due to the changes in use of factors in production. The following output decomposition model was used in this study. The Cobb-Douglas production function in logarithmic form for LLL method of wheat production is:

$$\ln Y_{LLL} = \ln b_0 + b_1 \ln X_{1LLL} + b_2 \ln X_{2LLL} + b_3 \ln X_{3LLL} + b_4 \ln X_{4LLL} + b_5 \ln X_{5LLL} + b_6 \ln X_{6LLL} + U_{LL} \quad \dots (1)$$

Where,

Y=Output of wheat and paddy (qtl);

X_1 = Human Labour (man days);

X_2 = Machine labour (hrs);

X_3 = Seed (kg);

X_4 = Fertilizer (kg);

X_6 = Plant Protection Chemicals (gm);

X_6 = Irrigation (hrs);

U = Random disturbance term, in conformity with the OLS assumptions;

b_0 = Scale parameter, and

b_i =slope parameters of the regression function (Production elasticities, $i=1$ to 6)

The per hectare production function for conventional tillage method is given in Equation as below:

$$\ln Y_{CT} = \ln a_0 + a_1 \ln X_{1CT} + a_2 \ln X_{2CT} + a_3 \ln X_{3CT} + a_4 \ln X_{4CT} + a_5 \ln X_{5CT} + a_6 \ln X_{6CT} + U_{CT} \quad \dots (2)$$

Taking difference between the equation (1) and (2), the following decomposition model was arrived as follows:

$$\ln [Y_{LLL} / Y_{CT}] = \{ \ln (b_0/a_0) \} + \{ (b_1-a_1) \ln X_{1CT} + (b_2-a_2) \ln X_{2CT} + (b_3-a_3) \ln X_{3CT} + (b_4-a_4) \ln X_{4CT} + (b_5-a_5) \ln X_{5CT} + (b_6-a_6) \ln X_{6CT} \} + \{ b_1 \ln (X_{1LLL} / X_{1CT}) + b_2 \ln (X_{2LLL} / X_{2CT}) + b_3 \ln (X_{3LLL} / X_{3CT}) + b_4 \ln (X_{4LLL} / X_{4CT}) + b_5 \ln (X_{5LLL} / X_{5CT}) + b_6 \ln (X_{6LLL} / X_{6CT}) \} + \{ U_{LLL} - U_{CT} \} \quad \dots (3)$$

The left hand side of the equation (3) gives the total difference in productivity expressed as an approximate percentage over conventional tilled farm. The natural logarithm of the ratio of per hectare net returns of LLL adopter farms and conventional tilled farms is approximately a measure of percentage difference in net outputs of the two categories of farmers. The first bracketed term on the right hand side, the natural logarithm of constant terms, is the gap attributable to the neutral component of technology. It is a measure of neutral technological gap. The second bracketed term is the gap attributable to the non-neutral component of technology weighted by input use expenditure for conventional tilled farms. That is, it is a measure of non-neutral technological gap, after adjustment in the level of input use expenditure weighted by the slope coefficients of the production function fitted for the LLL technology adopter farms. Hence, it is the gap due to difference in the levels of input use between two category of farmers after making due adjustment for production elasticities of different inputs. The last component is a random term which the model could not take into account.

The incremental benefit generated over conventional tillage farm under LLL is computed on per hectare and per LLL operation by equation (Aryal *et al.*, 2014) as follows:

$$\text{Incremental Benefit (Rs/LLL)} = \sum_{i=1}^n [P_{W_i} \Delta Y_{W_i} + P_{R_i} \Delta Y_{R_i}] - C_L H_L$$

Where,

P_{W_i} = Price of wheat per unit; ΔY_{W_i} = Additional yield of wheat in LLL field over CT farm

P_{R_i} = Price of rice per unit; ΔY_{R_i} = Additional yield of rice in LLL field over CT farm

C_L = Cost of hiring laser leveling service per hour; H_L = Time required to laser level the land

$i=1, \dots, n$ i.e. frequency of LLL operation (years)

Results and Discussion

In the study area, crop production was the major activity reported by more than 80 per cent of the sample farms (Table 1). The rice (*Oryza sativa*) crop was sown during the *kharif*

season (June to November), whereas wheat (*Triticum aestivum*) was the major crops grown extensively by the farmers in the *rabi* season (November to May).

Age of CSV farmers was lower than Non-CSV farmers. It is observed that younger people are more receptive to new technology. The family size of CSV farmers is smaller, are better educated and higher proportion had farming as a main occupation than those of Non-CSV farmers in study area. More number of CSV farmers were members of various organisations likes Farmers Club, Self-Help Groups, Grampanchayat, Co-operative societies etc. as compared to Non-CSV farmers, which could play a role in to get more exposure to the availability of various adaptation strategies to climate change. The CSV farmers have on an average three times more access to credit and five times more access to training as compared to Non-CSV farmers. The average farm size was also observed to be more in case of CSV farms (1.83 ha) as compared to Non-CSV farms (1.78 ha). At overall level, the cropping intensity on CSV sample farms is 18 per cent more than Non-CSV sampled farms.

Table 1: Table 4.1 Socio-economic characteristics of sample farm households

Sr. No.	Particulars	Vaishali farmers		Overall
		CSV	NCSV	
1	Sample size (No.)	39	40	79
2	Average age (years)	46.44	49.75	48.09
3	Average family size (No.)	7.4	7.7	7.55
4	Literacy (%)	65	57.5	61.25
5	Farming main occupation (%)	86.25	82.5	84.38
6	Farm size (ha)	1.83	1.78	1.81
7	Training access (Yes=1; otherwise=0)	61	7	34
8	Membership of organization (Yes=1; otherwise=0)	37	14	25.5
9	Credit access (Yes=1; otherwise=0)	56	19	37.5
10	Cropping intensity	176	158	167

Resource-use, Cost and Return Structure in Wheat and Paddy Production

The major farm inputs used for the production of wheat and paddy in LLL and CT methods are mentioned in Table 2.

Table 2: Major farm inputs used in wheat and paddy production in North-Eastern IGP (Per hectare)

Particulars	Laser Land Levelling	Conventional Tillage	Change (%)
Wheat			
Human labour (man days)	45	49	-10
Machine labour (hours)	12	11	6
Seeds (kg)	112	118	-5
Fertilizer (kg)	311	320	-3
PPC (gm)	312	356	-12
Duration of Irrigation (hr)	22	26	-17
Yield (Qtl)	40	37.6	6.38
Paddy			
Human Labour (man days)	71	76	-7
Machine Labour (hr)	11	10	7
Seed (Kg)	24	27	-11
Fertilizers (Kg)	252	264	-5
PPC (gm)	372	390	-5
Duration of Irrigation(hr)	53	65	-19
Yield (Qtl)	45.1	42.4	6.37

From the table it is revealed that for the wheat crop on an average about 10 per cent less human labour is required under

LLL farms. But the machine labour requirement was observed to be 6.19 per cent higher in case of LLL farms. A reduction in use of seed, fertilizers and PPC was observed to the extent of 5.08, 2.81 and 12.27 per cent, respectively under the LLL farms as compared to CT. The maximum reduction of seed was found on large farms, while that of fertilizers and PPC was observed on small farms. At overall farm size level, about 17 per cent reduction in duration of irrigation was observed in wheat crop under LLL. But as per farm wise reduction was more on large farms i.e. about 21.29 per cent.

While in case of paddy, LLL technology led to higher reduction in duration of irrigation as compared to that observed under wheat crop and it was 12 hours per hectare per season. The reduction was high on large farms. The use of human labour, seed, fertilizers and PPC was lower by about 6.58, 11.11, 4.55 and 4.62 per cent, respectively under LLL farmers compared to CT farms. The machine labour requirement was reported to have increased on LLL fields. Therefore, due to adoption of LLL technology about 16 hours of irrigation was saved in wheat and paddy cultivation. LLL also saved the input use especially human labour, seed, fertilizers and PPC as compared to that in CT and also increase in yield by 6.38 and 6.37 per cent, respectively was observed in the wheat and paddy crop.

The production costs and returns of wheat and paddy production using LLL and CT methods are presented in Table 3.

Table 3: Cost and return in wheat and paddy production using LLL method in NE IGP (Rupees per hectare)

Particulars	Laser Land Levelling	Conventional Tillage	Change (%)
Wheat			
Cost on human Labour	7788	8610	-10
Cost on machine Labour	7200	6780	6
Cost on seeds	3360	3540	-5
Cost on fertilizer	3888	4034	-4
Cost on PPC	1483	1690	-12
Irrigation charges	1080	1296	-17
Overhead Cost	2520	2595	-3
Total Operational cost	27318	28545	-4
Gross income	60400	56776	6
Return over operational cost	33082	28231	17
Benefit-cost ratio over cost C ₁	1.62	1.48	10
Cost of grain production (Rs/kg)	6.83	7.58	-10
Paddy			
Cost on human Labour	12780	13680	-7
Cost on machine Labour	6480	6072	7
Cost on seeds	768	864	-11
Cost on fertilizer	3175	3326	-5
Cost on PPC	1767	1853	-5
Irrigation charges	2628	3248	-19
Overhead Cost	2760	2904	-5
Total Operational cost	30358	31947	-5
Gross income	56375	53000	6
Return over operational cost	26017	21053	24
Benefit-cost ratio over cost C ₁	1.37	1.24	10
Cost of grain production (Rs/kg)	6.73	7.53	-11

From the table it is revealed that the gross returns of wheat production were Rs. 60400 per hectare in LLL and 56776 per ha in CT which was found to be 6 per cent more under LLL. The return over operational cost amounted to Rs. 33082 per ha in LLL and Rs. 28231 per ha in CT method of wheat production. The net income was higher in LLL method due to

higher yield and reduction in cost of cultivation as compared to CT method of wheat cultivation. The cost of cultivation amounted to Rs. 27318 per ha in LLL method and Rs.28545 per ha in CT method. The lower cost of cultivation was due to lower expenses on human labour (10 per cent), irrigation (17 per cent) and other inputs like seed, fertilizers and PPC in LLL than in CT method. The benefit-cost ratio of 1.62 was observed in LLL as against 1.48 in CT method of wheat production. The cost incurred to produce a kilogram of wheat was Rs. 7.58 in CT and Rs. 6.83 in LLL methods. Thus, the cost of wheat grain production was lower by 10 per cent in LLL as compared to in CT method.

However, in case of paddy production the total operational cost under LLL technology adopters' field was found to be reduced by 5 per cent over conventional tilled farm. In which the more reduction was observed in case of irrigation charges mainly due to reduction in duration of irrigation under LLL field. In all the paddy production under LLL helped to raise the return over operational cost by 24 per cent mainly due to reduction in cost incurred on inputs as well as due to incremental gain in productivity as compared to conventional tilled farm. The benefit-cost ratio was observed 10 per cent more in LLL as against CT method of paddy production. In all, significant difference was observed in wheat and paddy yield with and without LLL method. The cost incurred to produce a kilogram of paddy was Rs. 6.73 in LLL and Rs. 7.53 in CT methods. It indicates 11 per cent less under LLL paddy cultivation. This analysis suggests that LLL technology offers ample scope to generate additional income and help to conserve scarce resources. In this context, it is essential to assess the extent of change in productivity of wheat and paddy crop due to adoption of LLL technology and change in input use. Therefore, the productivity difference among adopters and non-adopters of laser land leveling for wheat and paddy production was segregated into its constituent sources i.e. technological change and input use differential with the help of decomposition model as suggested by Bisalial (1977) [8].

Decomposition Analysis

Using Equation (3), the values of production parameters and the geometric mean of input levels the total change in wheat and paddy output with the adoption of LLL technology was decomposed. Per hectare production of wheat and paddy was about 8 per cent and 6 per cent higher with LLL technology than with CT method. How much of this increased output was due to technological change and how much of it was due to change in input levels were also computed and are given in Table 4.

Wheat: The estimated improvement in productivity (per cent) with adoption of LLL technology was partly attributed to technological change and partly to input use, however, the two together contributed to 7.81 per cent of change in yield of wheat crop. However, the contribution of technology in yield increase was around 17 per cent. It indicates that the farmer could increase returns from wheat by 17 per cent just by shifting from conventional tillage to laser land levelling in NE IGP of India. As the laser land levelling technology has resource conserving potential, the impact of majority of inputs was observed to be negative in study area. The change in machine labour and fertilizers contributed to around 0.32 and 3.19 per cent of the increased output. Since, LLL is a resource conserving technology and gives more yield than CT it is

recommended to increase the existing productivity of wheat by increasing the level of adoption of LLL technology.

Table 4: Decomposition of Total Productivity Differences between LLL and CT Technology of Wheat and paddy in North-Eastern IGP. (Percent Contribution)

Sources of productivity Difference	Wheat	Paddy
Total observed difference in productivity	7.84	6.19
A. Due to difference in technology	17.48	46.63
B. Due to difference in input use level		
a) Human Labour	-0.75	-28.20
b) Machine Labour	0.32	0.94
c) Seed	-1.36	0.24
d) Fertilizer	3.19	2.17
e) Plant Protection Chemicals	-1.31	0.53
f) Irrigation	-9.75	-16.02
C. Total Estimated difference in productivity due to all resources (A+B)	7.81	6.28

Paddy

In case of paddy cultivation, the estimated improvement in productivity with LLL technology are due to technological change and due to input use and the two together technologies together contributed 6.28 per cent of total increase in yield. The estimated contribution of technology in increase in yield of paddy was 46 per cent. It indicates that the farmer could increase the yield of paddy crop by 46 per cent just by shifting from conventional tillage to laser land levelling. The impact of input use levels was observed to be negative mainly because of resource conserving potential of technology. Some inputs namely; machine labour, seed, fertilizers and PPC contributed around 0.94, 0.24, 2.17 and 0.53 per cent of the increased output. Hence in order to increase the existing productivity of paddy, there is need to increase in level of LLL adoption. Technology influences the sources of output growth by shifting the values of scale and slope parameters of the production function (Bisalial, 1977) [8].

Farmers' Perception on Impact of Laser Land Levelling Technology

In case of study area the adoption pattern of LLL technology was found more skewed towards medium and large farmers. Hence there is greater scope to extend the technology to the small farms. Majority of the farmers felt that there is decrease in time taken to irrigate the crop and decrease in water requirement; while some farmers reported that time required for field preparation remains same in LLL. The average time for leveling of one acre of land was 2 to 2.5 hours but it ranged from 2 hour to 3 hours depending on topography of land. All the farmers recorded water saving to a tune of 15-20 per cent after adoption of laser land leveler in wheat and rice crop. Most of the farmers recorded 6-8 per cent more yield under laser levelled field. Instead of number of advantages of LLL technology it was observed that the LLL technology was adapted to a limited extent in study area. There are number of constraints which restrict its adoption. The main reasons for non-adoption of LLL in Bihar was observed as small size of land holding followed by lack of technical knowledge and lack of awareness which restrict the adoption of technology. The Non-CSV farmers in study area are also facing the problem of poor institutional support which limits their exposure to LLL advantages. So there is need to increase the institutional support and provide technical knowledge to the farmers to enhance the adoption of LLL.

Benefits of using laser land levelling

When a farmer levels 1 ha of land using laser land leveller, the effect typically lasts for at least 3 years in NE IGP i.e. Bihar. Thus, it is assumed that the life of one levelling to be 3 years. Secondly it is also assuming that the farmer practices the dominant cropping pattern of rice-wheat during the entire period; planting rice in *kharif* and wheat in *rabi* season. Finally it is assumed that the increase in price of wheat and rice and the discount rate over time will balance each other so that prices stay constant at the current minimum support prices (MSPs) as fixed by the government of India. Based on above mentioned assumptions the benefit derived from LLL operation is compiled and presented in Table 5. The incremental benefit to the farm due to use of LLL technology is calculated by using the equation as given below:

$$\text{Incremental Benefit (Rs. / LLL)} = \sum_{i=1}^3 [P_{wi}\Delta Y_{wi} + P_{ri}\Delta Y_{ri}] - C_L H_L$$

Table 5: Incremental benefit of using LLL

Sr. No.	Particulars	Incremental Benefit
1	Price of Wheat (Rs/kg)	16.25
2	Price of paddy (Rs/kg)	15.5
3	Incremental yield of Wheat(kg/ha)	240
4	Incremental yield of paddy(kg/ha)	270
5	Benefit per year from wheat(Rs/ha)	3900
6	Benefit per year from paddy (Rs/ha)	4185
7	Number of years LLL applicable	3
8	Benefit per LLL operation	24255
9	Cost of LLL (Rs/ha)	4200
10	Net Benefit per year (Rs)	6685
11	Net Benefit per LLL operation	20055

The benefit stream is calculated by using minimum support price (2017-18) for respective crop which is Rs.16.25 per kg for wheat and Rs.15.50 per kg for rice. The study consider the yield differential in laser leveled land and conventional tilled land to be 240 kg/ha for wheat and 270 kg/ha for in rice. Based on farmers' response, in study area the cost of hiring laser land leveling service was Rs. 700 per hour and on an average around 6 hours time was required to level a hectare of land by LLL technology. Therefore, the total cost of levelling one hectare is Rs. 4200. In all, about Rs.20055 of net incremental benefit was derived from the adoption of LLL in study area.

Conclusions

The study has revealed that it is possible to save scarce resources and irrigation water under LLL than under conventional method. Due to resource saving, net return has been significantly higher in LLL technology. Hence, this technology is an important alternative to save scarce resources and enhance the net farm income. The decomposition analysis has shown that per hectare production of wheat and paddy was 7.81 per cent and 6.28 per cent, respectively higher in LLL than in conventional tillage method. In this improved production method, LLL technology contributed around 17 per cent and 46 per cent change in productivity of wheat and paddy crops, respectively. By adopting this technology, farmers could save scarce resources and reduce the cultivation cost. The constraint of small size of land holding can be solved by co-operative or collective farming. The availability of institutional support and provision of technical knowledge

to the farmers needs to be accorded more attention to foster the adoption of LLL technology.

References

1. Abdullaev I, Hasan MU, Jumaboev K. Water saving and economic impacts of land leveling: the case study of cotton production in Tajikistan. *Irrigation Drainage System*. 2007; 21:251-263.
2. Aggarwal PK, Joshi PK, Ingram JS, Gupta RK. Adapting food systems of the Indo-Gangetic plains to global environmental change: key information needs to improve policy formulation. *Environmental Science & Policy*. 2004; 7(6):487-498.
3. Aggarwal PK, Singh AK, Samra JS, Singh G, Gogoi AK, Rao GGSN, Ramakrishna YS. Introduction in global climate change and Indian agriculture, Ed: PK Aggarwal, ICAR, New Delhi, 2009, 1-5.
4. Ahmed B, Khokhar SB, Badar H. Economics of laser land leveling in district Faisalabad. *Pakistan Journal of Applied Science*. 2001; 1(3):409-412.
5. Aryal JP, Bhatia M, Jat ML, Sidhu HS. Impacts of laser land leveling in rice-wheat rotations of the North-western Indo-Gangetic Plains of India. In *World Congress of Environmental and Resource Economics*, 2014.
6. Aryal JP, Mehrotra MB, Jat ML, Sidhu HS. Impacts of laser land leveling in rice-wheat systems of the north-western Indo-gangetic plains of India. *Food Security*. 2015; 7(3):725-738.
7. BIRTHAL PS, KHAN MT, NEGI DS, AGARWAL S. Impact of climate change on yields of major food crops in India: Implications for food security. *Agricultural Economics Research Review*. 2014; 27(2):145-155.
8. Bisalial S. Decomposition Analysis of Output Change under New Production Technology in Wheat Farming: Some Implications to Returns on Research Investment, *Indian Journal of Agricultural Economic*. 1977; 32(3):193-201.
9. Branca G, McCarthy N, Lipper L, Jolejole MC. Climate-smart agriculture: a synthesis of empirical evidence of food security and mitigation benefits from improved cropland management. *Mitigation of Climate Change in Agriculture Series*. 2011; 3:1-42.
10. FAO. Climate Smart Agriculture: Policies, Practices and Financing for Food Security, Adaptation and Mitigation. Food and Agriculture Organization of the United Nations (FAO), Rome, Italy. Retrieved from, 2010. <http://www.sidalc.net/cgi-bin/wxis.exe/?IsisScript=INTER.xis&method=post&formato=2&cantidad=1&expression=mfn=000713>.
11. Gill GJ. An assessment of the impact of laser-assisted precision land levelling technology as a component of climate-smart agriculture in the state of Haryana, India. New Delhi: CIMMYT-CCAFS, International Maize and Wheat Improvement Center (CIMMYT), 2014.
12. Gupta RK, Gill MA. New opportunities for saving on water: Role of resource conserving technologies. Addressing resource conservation issues in rice-wheat systems of South Asia (No. CIS-3599. CIMMYT), 2003.
13. Jat ML. Laser land leveling in India: A success. In *Lessons Learned from Postharvest and Mechanization Projects, and Ways Forward*. Los Banos, Philippines, May, 2012, 22-24.
14. Jat ML, Chandna P, Gupta R, Sharma SK, Gill MA. Laser land leveling: A precursor technology for resource

conservation. Rice-Wheat consortium technical bulletin series. 2006; 7:48.

15. Jat ML, Gupta R, Saharawat YS, Khosla R. Layering precision land leveling and furrow irrigated raised bed planting: Productivity and input use efficiency of irrigated bread wheat in Indo-Gangetic Plains. *American Journal of Plant Sciences*. 2011; 2(04):578.
16. Jat ML, Sharma SK, Gupta RK, Sirohi K, Chandana P. Laser land leveling: the precursor technology for resource conservation in irrigated ecosystem of India. I: Abrol I P, Gupta RK and Malik RK (ed) *Conservation Agriculture-status and prospects CASA*, New Delhi, 2005, 145-154.
17. Joshi PK, Tyagi NK. Salt affected and water logged soils in India: a review. *Strategic Change in Indian Irrigation*. ICAR, New Delhi, India, and IFPRI, Washington, DC, USA, 1994, 237-252.
18. Kahlowan MA, Raouf A, Hanif M. Rice yield as affected by plant densities. *Mona Reclamation Experimental Project Bhalwal (238)*. Report, 2000.
19. Kerr RA. Northern India's Groundwater Is Going, Going, Going.... *Science*, 2009; 325(5942):798.
20. Kumar V, Kumar D, Kumar P. Declining water table scenario in Haryana-A review. *Water and Energy International*. 2007; 64(2):32-34.
21. Thakar S, Gill MS, Sidhu HS, Sukhpreet S, Sidhu AS. Effect of laser land leveling on water productivity and yield of rice-wheat cropping system. *Indian Journal of Ecology*. 2009; 36(1):95-97.