



ISSN 2278- 4136

Online Available at [www.phytojournal.com](http://www.phytojournal.com)

## Journal of Pharmacognosy and Phytochemistry

# Preliminary Phytochemical Analysis of the Extracts of *Psidium* Leaves

Vikrant Arya<sup>1\*</sup>, Narender Thakur<sup>2</sup>, C.P. Kashyap<sup>3</sup>

1. College of Ayurvedic Pharmaceutical Sciences, Jogindernagar, Mandi H.P., India.  
[E-mail: [arya.vikrant30@gmail.com](mailto:arya.vikrant30@gmail.com)]
2. Vinayaka College of Pharmacy, Village Bahuguna P.O. Garsa Distt. Kullu, H.P., India
3. College of Ayurvedic Pharmaceutical Sciences, Jogindernagar, Mandi H.P., India.

---

Amritphale (*Psidium guajava* L.), is a small tree in Myrtaceae family, traditionally used in treatment of several diseases (inflammation, diabetes, hypertension, wounds, pain and fever). The present study was carried out to investigate the phytochemical profile of leaves of *Psidium guajava* L. The leaves powder was successively extracted with petroleum ether, chloroform, ethanol, water, hydroalcoholic. Phytochemical analysis shows the presence of flavonoids, tannins triterpenoids, saponins, sterols, alkaloids and carbohydrates. The result of the study could be useful for description and foundation of monograph of the plant.

---

**Keyword:** Flavonoids, Psidium, Phytochemical

### 1. Introduction

Quality can be defined as the status of a drug that is determined by identity, purity, content and other chemical, physical, or biological properties, or by the manufacturing processes. Quality control is a term that refers to processes involved in maintaining the quality and validity of a manufactured product. For the quality control of a traditional medicine, the traditional methods are procured and studied, and documents and the traditional information about the identity and quality assessment are interpreted in terms of modern assessment. *Psidium guajava* is a small tree upto 20 feet high, with spreading branches. Guava is easy to recognize because of its smooth, thin, copper-colored bark that flakes off, showing the greenish layer beneath and also because of the attractive, "bony" aspect of its trunk which may in time attain a diameter of 10 inch<sup>[1]</sup>. *Psidium guajava*, which is considered a native to Mexico,

extends throughout the South America, European, Africa and Asia. It is mainly found in tropical and subtropical regions. In India, it is distributed in Uttar Pradesh, Bihar, Maharashtra, Assam, West Bengal, Haryana and Andhra Pradesh<sup>[2]</sup>.

The leaves are evergreen, opposite, short-petioled, oval or oblong-elliptic, somewhat irregular in outline; 7-15 cm long to 3-5 cm wide, leathery, with conspicuous parallel veins, and more or less downy on the underside<sup>[3]</sup>. Flowers are white, borne singly or in small clusters in the leaf axils. Flowers are 2-3 cm wide, with 4 or 5 white petals which are quickly shed, and a prominent tuft of perhaps 250 white stamens tipped with pale-yellow anthers<sup>[4]</sup>. The fruit, exuding a strong, sweet, musky odor when ripe, may be round, ovoid, or pear-shaped, 2.5-10 cm long, with 4 or 5 protruding floral remnants (sepals) at the apex with thin, light-yellow skin, frequently blushed with pink. Next to the skin is a

layer of somewhat granular flesh, 1/8 to 1/2 inch thick, white, yellowish, light- or dark-pink, or near-red, juicy, acid, subacid, or sweet and flavorful. The central pulp, slightly darker in tone, is juicy and normally filled with very hard, yellowish seeds, 1/8 inch long, though some rare types have soft, chewable seeds. Actual seed counts have ranged from 112 to 535 but some guavas are seedless or nearly so. When immature and until a very short time before ripening, the fruit is green, hard, gummy within and very astringent. Bark is quite smooth, pale pinkish brown usually tinged with chlorophyll [5].

### Taxonomic Classification

<i>Kingdom</i>	<i>Plantae</i>
<i>Subkingdom</i>	<i>Tracheobionta</i>
<i>Superdivision</i>	<i>Spermatophyta</i>
<i>Division</i>	<i>Magnoliophyta</i>
<i>Class</i>	<i>Magnoliopsida</i>
<i>Subclass</i>	<i>Rosidae</i>
<i>Order</i>	<i>Myrtales</i>
<i>Family</i>	<i>Myrtaceae</i>
<i>Genus</i>	<i>Psidium L.</i>
<i>Species</i>	<i>Guajava L.</i>

More recent ethnopharmacological studies show that *Psidium guajava* is used in many parts of the world for the treatment of a number of diseases, e.g. as an anti-inflammatory, for diabetes, hypertension, caries, wounds, pain relief and reducing fever. Some of the countries with a long history of traditional medicinal use of guava include Mexico and other Central American countries including the Caribbean, Africa and Asia [6].

## 2. Materials and method:


### 2.1 Collection of Plant Materials and Leaves:

The leaves were collected from fields of Kangra, Himachal Pradesh, India. The collected leaves were shade dried for 35 days and finally pulverized in to coarse powder. It was stored in a well closed container free from environmental climatic changes till usage.

### 2.2 Preparation of Extracts:

Successive extraction of plant material (Fig.1)

**Fig 1:** A Schematic Representation of Successive extraction



### 2.3 Preparation of Hydroalcoholic Extract

Hydroalcoholic (8:2) extract was prepared using leaves of *Psidium guajava*. Dried leaves (250 gm) were ground. The ground sample was soaked in hydroalcoholic solvent (8:2 v/v) and left for 24 h. The mixture was filtered and the filtrate concentrated by evaporation at 40 °C. Then extract was dried and weighed.

### 2.4 Preliminary Phytochemical Screening

The preliminary phytochemical tests were performed for testing different chemical groups present in extracts [7-8]. Extract (100 mg) was treated with few drops of Dragendorff's reagent [Potassium bismuth iodide solution]. Formation of orange brown precipitate indicated the presence of alkaloids. To 100 mg of extract small quantity of Wagner's reagent [Solution of iodine in potassium iodide] was added. Presence of reddish brown precipitate if alkaloids are present. To 100 mg of extract small quantity of Hager's

reagent [saturated solution of Picric acid] was added. Formation of yellow precipitate indicated the presence of alkaloids. Powdered drug extract on shaking vigorously with water results into persistent foam. 200 mg extract was boiled with 3 mL of dil.  $H_2SO_4$  in a test tube for 5 min and filtered while hot. Cool and added the equal volume of  $C_6H_6$  and  $CHCl_3$ , shake well and separated the organic solvent and added the  $NH_3$ . The ammonical layer turned pink or red. Alcoholic extract was treated with 1 mL pyridine and 1 mL of sodium nitroprusside. Pink to red colour appears. Extract (2 mL) was treated with 0.4 mL of glacial acetic acid containing a trace amount of  $FeCl_3$  and 0.5 mL of concentrated  $H_2SO_4$  was also added by the side of the test tube. Persistent blue color appeared in the acetic acid layer if cardiac glycosides were present. To 5 mL of extract few drops of 5%  $FeCl_3$  was added. Presence of deep blue black colour indicated the presence of tannins. When 5 mL of extract was treated with few drops of 5% lead acetate solution, white precipitates appeared. To 5 mL of extract 5 mL of 95% ethanol was added along with dilute HCl from sides of test tube. Few fragments (0.5 g) of magnesium turnings were also added. Presence of slight pink colour indicated the presence of flavonoids. To 5 mL of extract few drops of NaOH solution was added. Formation of an intense yellow color, which turns to colorless on addition of few drops of dil.  $H_2SO_4$  indicated the presence of flavonoids. A little extract was taken with 2 mL of water and 0.5 mL of concentrated  $HNO_3$  was added to it. Yellow colour is obtained if proteins are present. To 5 mL of extract 4% NaOH was added along with few drops of 5%  $CuSO_4$  solution. Violet or pink colour appeared indicated the presence of proteins. Extract (5 mL) was treated with 5 mL  $CHCl_3$  with few drops of conc.  $H_2SO_4$ , shake well and allowed to stand for some time. Formation of yellow colored lower layer indicated the presence of triterpenoids. Extract (5 mL) was treated with few drops of acetic anhydride, boiled and cooled, conc.  $H_2SO_4$  was added from the sides of the test tube showed a brown ring at the junction of two layers and the upper layer turns green which showed the presence of Steroids and formation of

deep red color indicated the presence of triterpenoids. In a test tube containing 5 mL of extract, few drops of freshly prepared 10% alcoholic solution of  $\alpha$ -naphthol was added and shaken/stirred for few min. Then 5 mL of conc.  $H_2SO_4$  was added from sides of the test tube. Violet ring was formed at the junction of two liquids, indicated the presence of carbohydrates. Small quantity of extract was pressed the between two filter papers, the stain on 1<sup>st</sup> filter paper indicated the presence of fixed oils. The extract was evaporated to get 10 mL of extract. To the extract 25 mL of 10% NaOH was added, then it was boiled in water bath for 30 min. The extract was cooled and excess of sodium sulphate was added. Soap was formed at the top and filtered. To the filtrate  $H_2SO_4$  was added which was evaporated. The extract was dissolved in ethanol and few drops of  $CuSO_4$  and NaOH was added. Clear blue solution indicated the presence of fats.

**Figure : Fruits, Flowers, Leaves and Seeds of *Psidium guajava* Linn**



### Observations:

**Table 1:** Successive and Hydroalcoholic Extraction of Powder of *Psidium guajava* Linn. Leaf

Type of extract	Amount of extract (gm)	Yield (% w/w)	Appearance
Petroleum ether	2.850	1.54	Oily yellowish black
Chloroform	10.175	4.47	Greenish Brown
Ethanollic	18.013	7.61	Brownish black solid mass
Aqueous	17.650	7.46	Brown
Hydroalcoholic	20.425	8.57	Brown

**Table 2:** Phytochemical Screening of Various Extracts of *Psidium guajava* Linn.

Chemical tests	Petroleum ether extract	Chloroform extract	Ethanol extract	Aqueous extract	Hydroalcoholic extract
<b>Alkaloids</b>					
<i>Mayer's reagent</i>	-	-	-	-	-
<i>Dragendorff's reagent</i>	-	-	+++	-	-
<i>Wagner's reagent</i>	-	-	-	-	-
<i>Hager's reagent</i>	-	-	++	-	++
<b>Saponins</b>					
<i>Froth test</i>	-	-	-	-	++
<b>Sterols</b>					
<i>Salkowski test</i>	-	-	++	-	++
<i>Leibermann's reagent</i>	-	-	++	++	++
<i>Leibermann Burchard's</i>	-	-	++	++	++
<b>Carbohydrates</b>					
<i>Molisch's test</i>	-	-	++	++	++
<i>Fehling's test</i>	-	-	++	++	++
<i>Camelisation</i>	-	-	++	++	++
<b>Anthraquinone glycosides</b>					
<i>Borntrager's test</i>	-	-	-	-	-
<b>Cardiac glycosides</b>					
<i>Legal's test</i>	-	-	-	-	-
<i>Keller-Killiani test</i>	-	-	-	-	-
<b>Tannins</b>					
<i>Lead acetate solution</i>	-	++	+++	+++	+++
<i>Ferric chloride solution</i>	-	-	++	++	++
<b>Proteins</b>					
<i>Xanthoproteic test</i>	-	-	-	-	-
<i>Biuret test</i>	-	-	-	-	-
<b>Flavonoids</b>					
<i>Ammonia test</i>	-	-	++	++	++
<i>Alkaline reagent test</i>	-	++	++	++	++
<i>Magnesium ribbon test</i>	-	-	++	++	++

(+) Positive Test, (-) Negative test

### 3. Conclusion

Phytochemical screening of petroleum ether, chloroform, ethanol, aqueous and hydroalcoholic extracts revealed the presence flavonoids, tannins triterpenoids, saponins, sterols, alkaloids and carbohydrates by positive reaction with the respective test reagent. Phytochemical screening showed that maximum presence of

phytoconstituents in ethanolic and hydroalcoholic extracts.

### 4. Acknowledgement:

The authors are grateful to Mr. N.S Thakur, Chairman Vinyaka College of Pharmacy and Dr. C.P. Kashyap, Principal, College of Ayurvedic Pharmaceutical Sciences Jogindernagar, Mandi,

H.P. for providing facilities to carrying out research on *Psidium* leaves.

## 5. Reference

1. Morton JF. Guava. In: Fruits of warm climates. Miami FL 1987; 356–363.
2. Kamath JV, Nair R, Kumar CKA. *Psidium guajava* L: A Review. Int J Green Pharma 2008; 2(1):0-12.
3. Yusof SG. University Putra Malaysia, Selangor, Malaysia. Elsevier Science Ltd 2003; 2985-2991.
4. Stone B. The flora of Guam. Micronesica Behav 1970; 82(2):373-378.
5. Morton JF. Guava. In: Fruits of warm climates. Miami FL 1987; 356–363.
6. Kirtikar KR, Basu BD. Indian Medicinal Plants. Vol. 1, International Book Distributors, Dehradun, 1998, 1045-1048.
7. Kokate CK. Practical Pharmacognosy. Vallabh prakashan, Delhi, 2005, 107-111.
8. Khandelwal KR. Practical Pharmacognosy Techniques and Experiments. Edn 9, Nirali Prakashan, Pune, 2003.