



E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2015; 3(6): 67-73
Received: 17-01-2015
Accepted: 01-02-2015

Abd ELnasser B. Sengab
Pharmacognosy Department,
Faculty of Pharmacy, Ain Shams
University, Cairo, Egypt.

Dina M. Y. El naggar
Pharmacognosy Department,
Faculty of Pharmacy, Al azhar
University, Cairo, Egypt.

Mohamed R. Elgindi
Pharmacognosy Department,
Faculty of Pharmacy, Egyptian
Russian University, Helwan
University, Cairo, Egypt.

Mostafa B. Elsaid
Pharmacognosy Department,
Faculty of Pharmacy, Egyptian
Russian University, Helwan
University, Cairo, Egypt.

Correspondence:

Mohamed R. Elgindi
Pharmacognosy Department,
Faculty of Pharmacy, Egyptian
Russian University, Helwan
University, Cairo, Egypt.

Biological studies of isolated triterpenoids and phenolic compounds identified from *Wodyetia bifurcata* family Areaceae

Abd ELnasser B. Sengab, Dina M. Y. El naggar, Mohamed R. Elgindi and Mostafa B. Elsaid

Abstract

The chromatographic investigation of the aerial parts of *Wodyetia bifurcata* (family Areaceae) revealed the presence of β -Amyrin, lupeol, apigenin, kaempferol, p- hydroxyl benzoic acid and gallic acid that isolated for first time from *Wodyetia bifurcata* and their structures are elucidated by different spectroscopic techniques as UV, $^1\text{H-NMR}$, $^{13}\text{C-NMR}$ and EI/MS spectra. The biological activities of different plant extracts are tested for anti-inflammatory, antioxidant and cytotoxicity.

Keywords: *Wodyetia bifurcata*, Areaceae, Aerial part, Triterpenoids, phenolic compounds, Anti-inflammatory, Antioxidant, Cytotoxicity.

1. Introduction

Wodyetia is a genus of trees belonging to family Areaceae. *Wodyetia bifurcata* L (also called foxtail) is fast growing, large sized evergreen tree, up to 10 m tall, grows in El-Orman garden, Giza, Egypt. *Wodyetia bifurcata* L is endemic to Australia and it is found on north Queensland [18]. Triterpenoids are prevalent in the plant kingdom. Recent evidences support the beneficial effects of naturally occurring triterpenoids against several types of human diseases [27], anticancer [19], antioxidant [2], hepatoprotective [22], antifungal [28], antibacterial [16] and antileishmanial activity [9]. Flavonoids are a group of polyphenolic compounds, diverse in chemical structure and characteristics, found ubiquitously in plants. Until now, more than 9000 different flavonoid compounds were described in plants, where they play important biological roles by affecting several developmental processes [37]. There has been increasing interest in the research of flavonoids from dietary sources, due to growing evidence of the versatile health benefits of flavonoids including antioxidant activity [10], coronary heart disease [35], hepatoprotective [34], anti-inflammatory, [23], anticancer activities [38], antibacterial [21] while some flavonoids exhibit potential antiviral activities [5].

2. Results and discussions

The MS of the amyryns is characterized by an M^+ of m/e 426 and a major fragmentation yielding a component of m/e 218 [15]. Data of $^1\text{H NMR}$ for compound I showed eight methyl group corresponding to terpenoid of *olean* skeleton. The $^{13}\text{C NMR}$ and EI-MS of compound I were compared with that reported in that literature review [7, 8, 25]. The compound I is identified as β -Amyrin.

The $^1\text{H NMR}$ data of compound II showed seven tertiary methyl singlets and one secondary hydroxyl group. It also showed olefinic protons at δ 4.41 and 4.67. The compound showed 30 signals for the terpenoid of lupine skeleton which is represented by seven methyl groups. The carbon bonded to the hydroxyl group C-3 appeared at δ 79.1, while the alkenic carbons appeared at δ 151.1 and 109.5. EI-MS spectrum of compound II showed the molecular ion at m/z 426 (10%) [M^+] corresponding to the formula $\text{C}_{30}\text{H}_{50}\text{O}$, together with fragments at m/z 411 (6%) [M^+-15] and 408 (5%) [M^+-18] which were due to the loss of methyl group and a molecule of water from the molecular ion peak other fragments occurring at m/z 385 (4%) (M^+-41), m/z 220 (8%) ($\text{M}^+-\text{C}_{15}\text{H}_{26}$), m/z 218 (46%) ($\text{M}^+-\text{C}_{14}\text{H}_{20}\text{O}$), m/z 207(51%) ($\text{M}^+-\text{C}_{16}\text{H}_{27}$) and m/z 189 (70%) ($\text{M}^+-\text{C}_{14}\text{H}_{21}$) were characteristic for lupane series [15]. Data of $^1\text{H NMR}$, $^{13}\text{C NMR}$ and EI-MS were compared with that reported in that literature review [1, 3, 13, 36]. The compound II is identified as lupeol.

UV spectral data of compound III showed two major absorption bands; band I at 331 nm and band II at 272 nm, which is typical of a flavone nucleus with no hydroxyl group at position 3^[20]. The addition of sodium methoxide resulted in a bathochromic shift (+ Δ 69 nm) in band I with an increase in intensity, which proved that position 4' has a free OH group^[20]. The bathochromic shift in band I (+ Δ 56 nm) on addition of AlCl₃; compared with the same band in MeOH, which is still stable even after the addition of HCl indicated the presence of free 5-OH group. The bathochromic shift in band II on addition of NaOAc (+ Δ 9 nm) compared with the same band in MeOH suggested the presence of a free hydroxyl group at carbon number 7. On addition of H₃BO₃ to NaOAc the hypochromic shift in band I (- Δ 45 nm) suggested the absence of any *ortho*-dihydroxyl groups. The ¹H-NMR spectrum of compound III revealed the presence of two aromatic protons of the A-ring revealed as two doublet at δ 6.2 and δ 6.0 each proton has $J=1.5$ Hz due to *meta* coupling assigned to H-8 and H-6, respectively. H-3 appeared at δ 6.4 as a single signal as no neighbor proton appears. The aromatic protons of the B-ring as two doublets at δ 7.6 and δ 7.0 each doublet with $J=5.7$ Hz due to *ortho* coupling assigned to H-2', 6' (H-2', 6' are superimposed) and H-3', 5', (H-3', 5' are superimposed) respectively. Note that the H-3', 5' doublet occurs upfield from H-2', 6' due to shielding effect of the hydroxyl group and the deshielding influence of C-ring on H-2', 6'. The EI-MS spectrum of compound III showed the molecular ion peak as the base peak (M⁺) at m/z 270 (95%) corresponding to molecular formula C₁₅H₁₀O₅. Peaks at m/z 152 (43%) and m/z 118 (48%) it suggests the fragments of ring (A) and ring (B). Others peaks suggests another pathway of decomposition of compound and splitting of ring (A) and ring (B) at m/z 121 (20%) and m/z 149 (26%). Through the UV, EI-MS and ¹H-NMR the compound was identified as apigenin (isolated for the first time from *Wodeytia bifurcata*). Data of ¹H NMR, UV spectra and EI-MS were compared with that reported in that literature review (Guang-ying Chen, *et al.*, 2010; Fairouz Moussaoui *et al.*, 2010; Venkata Sai Prakash Chaturvedula and Indra Prakash, 2013; Jie Zhanga, *et al.*, 2005). The compound III is identified as.

UV spectral data of compound IV in methanol showed two major absorption bands; band I at 367 nm and band II at 265 nm, which is typical of a flavanol^[20]. The addition of sodium methoxide resulted in a bathochromic shift (+ Δ 53 nm) in band I with an no decrease in intensity, which proved that position 4' has a free OH group^[20]. The bathochromic shift in band I (+ Δ 47 nm) on addition of AlCl₃; compared with the same band in MeOH, which is still stable even after the addition of HCl indicated the presence of free 5-OH group. The bathochromic shift in band II on addition of NaOAc (+ Δ 11 nm) compared with the same band in MeOH suggested the presence of a free hydroxyl group at carbon number 7 and in band I there is lesser than NaOMe indicate that there is no substitution of hydroxyl group number 7. On addition of H₃BO₃ to NaOAc the hypochromic shift in band I (- Δ 15 nm) suggested the absence of any *ortho*-dihydroxyl groups. ¹H-NMR spectrum of compound IV showed the two aromatic protons of the A-ring revealed as two doublet at δ 6.5 and δ 6.2 each proton has $J=1.2$ Hz due to *meta* coupling assigned to H-8 and H-6, respectively. The aromatic protons of the B-ring as two doublets at δ 7.94 and δ 6.9 each doublet with $J=5.4$ Hz due to *ortho* coupling assigned to H-2', 6' (H-2', 6' are superimposed) and H-3', 5', (H-3', 5' are superimposed)

respectively. Note that the H-3', 5' doublet occurs upfield from H-2', 6' due to shielding effect of the hydroxyl group and the deshielding influence of C-ring on H-2', 6'. The EI-MS spectrum of compound IV showed the molecular ion peak as the base peak (M⁺) at m/z 287 (84%) corresponding to molecular formula C₁₅H₁₁O₆. Spectra showed peak at 269 m/z indicate that molecular formula C₁₅H₆O₅ (M-18) and removing of H₂O due to flavanol. It showed peak at m/z 259 (12%) suggest of (M- CO) and m/z 258 (15%) of (M- CHO). Peaks at m/z 153 (42%) and m/z 133 (48%) it suggests separation of ring (A) and ring (B). Others peaks suggests another pathway of decomposition of compound and splitting of ring (A) and ring (B) at m/z 121 (8%) and m/z 165 (%14). Through the UV, EI-MS and ¹H-NMR compound (F2) was identified as kaempferol (was isolated for the first time from *Wodeytia bifurcata*) data of ¹H NMR, UV spectra and EI-MS were compared with that reported in that literature review^[6, 24, 31]. The compound IV is identified as kaempferol.

Compound V was isolated from soluble part of successive methanol extract of *Wodeytia bifurcata* as Colorless crystals; it gave violet color with ferric chloride solution indicating the presence of phenolic group with melting point 215-216 °. The ¹H-NMR of compound V showed signals at δ ; 7.58 (2H, d, $J=2.1$ Hz, H-2, 6), 6.92 (2H, d, $J=1.2$ Hz, H-3, 5). Data of ¹H NMR, was compared with that reported in that literature review^[29]. The compound V is identified as para hydroxyl benzoic acid.

The compound VI is identified by co-chromatography with authentic using paper chromatography with solvent system butanol : acetic acid : water (5:4:1) and acetic acid 6%. Spots visualized under UV light show violet color and gave blue color with FeCl₃ with R_f = 0.53 and 0.56 respectively. The compound VI is identified as Gallic acid.

The biological experiments showed that methanol extract have potent inhibitory activity (82.1%) against LPS (Lipo-Poly-Saccharide) - induced nitric oxide to extent compared with that of dexamethasone. Butanol showed the least by (27.9%) and chloroform and n-hexane (42.6%) and (65.8%) respectively.

3. Experimental

3.1 Plant Material

The aerial parts of *Wodeytia bifurcata* were collected from the Egyptian Orman garden, Giza, Egypt at June 2011. The plant was identified by Agricultural Engineer Terese Labib, El Orman Botanical Garden.

3.2 Extraction and isolation

The air dried powdered leaves and stems of *Wodeytia bifurcata* (1 kg) were macerated in 3 L 80% aqueous methanol with occasional stirring at room temperature for three days. The methanolic extract was concentrated and dried under vacuum. The dried residue (85 g) was suspended with H₂O and extracted with butanol. The butanol extract was separated and evaporated to dryness under reduced pressure to yield a semisolid residue (Fraction I, 28 g). The aqueous liquor was evaporated till dryness under vacuum to yield a residue (Fraction II, 35 g).

The fraction I subjected to column silica gel achieved with gradient elution using chloroform: methanol (100% CHCl₃ to 100% MeOH) to yield (fraction I-a, 85:15 chloroform: methanol) and (fraction I-b, 80:20 chloroform: methanol) which are subjected to further purification by silica gel to yield compound I and compound II subjected to more purification

with preparative-TLC silica gel G F₂₄₅ eluate with CHCl₃: MeOH: H₂O (13:7:2).

The fraction II subjected to column silica gel achieved with gradient elution using chloroform: methanol (100% CHCl₃ to 100% MeOH) to yield (fraction II-c, 70:30 chloroform: methanol) and (fraction II-d, 40:60 chloroform: methanol) which are major fraction which is subjected to further purification by silica gel column to yield fractions which were further purified by rechromatography on sephadex LH-20 column using MeOH, MeOH/H₂O and H₂O as eluents to yield compound III, compound IV, compound V and compound VI.

Compound (I)

¹H NMR (CDCl₃, 300 MHz) at δ; 0.68, 0.78, 0.80, 0.86, 0.90, 0.93, 1.05, 1.18 (each 3H, s, C-25, C-23, C-30, C-29, C-24, C-26, C-27, C-28), 3.1(dd, *J*=7.2, 3.1 Hz, H-3) 5.7(d, *J*=7.2, Hz, H-12).

¹³C NMR (CDCl₃, 300 MHz): at δ; 15.12 (C-24), 15.12 (C-25), 16.82 (C-26), 18.77 (C-6), 20.53 (C-11), 23.15 (C-30), 23.44 (C-27), 24.06 (C-16), 25.69 (C-15), 26.54 (C-2), 27.48 (C-23), 28.43 (C-28), 30.12 (C-20), 31.42 (C-7), 33.21 (C-17), 33.78 (C-29), 34.42 (C-21), 35.25 (C-10), 35.94 (C-22), 37.25 (C-4), 37.40 (C-1), 39.32 (C-8), 41.42 (C-14), 46.12 (C-9), 48.42 (C-19), 48.70 (C-18), 55.12 (C-5), 79.32 (C-3), 122.91 (C-12), 145.66 (C-13). EI-MS spectrum: showed the molecular ion at *m/z* 426 (14%) [M⁺] corresponding to the formula C₃₀H₅₀O, together with fragments *m/z* 411(5%), 275 (3%), *m/z* 257 (2%), 218 (95%), 203 (46%), 189 (33%), 175 (17%), 161 (3%).

Compound (II)

¹H NMR (CDCl₃, 300 MHz) at δ; 0.77, 0.86, 0.88, 0.90, 0.92, 0.96, 1.11 (each 3H,s), 3.64 (1H, dd, *J*=5.2 Hz,H-3) 4.41 (1H,s,H-29a), 4.67 (1H,s,H-29b) showed seven tertiary methyl singlets and one secondary hydroxyl group. It also showed olefinic protons at δ 4.41 and 4.67. ¹³C NMR (CDCl₃,300MHz): at δ; 14.36 (C-27), 15.73 (C-24), 16.77 (C-26), 17.62 (C-25), 18.42 (C-28), 19.12 (C-6), 20.53 (C-30), 21.33 (C-11), 25.55 (C-2), 26.63 (C-12), 27.21 (C-15), 28.43 (C-23), 30.48 (C-21), 34.33 (C-7), 35.32 (C-16), 37.66 (C-10), 38.42 (C-1), 39.88 (C-4), 40.22 (C-13), 41.82 (C-22), 42.12 (C-8), 43.42 (C-14), 44.22 (C-17), 49.56 (C-19), 49.84 (C-18), 50.12 (C-9), 54.29 (C-5), 77.40 (C-3), 110.89 (C-29), 150.63 (C-20).

Compound (III)

UV spectral data at λ_{max} (nm): MeOH; 272, 331, NaOMe; 268, 332, 400, AlCl₃; 279, 306, 351, 387, AlCl₃/HCl; 279, 304, 346, 386, NaOAc; 281, 306, 387, NaOAc/H₃BO₃; 268, 342. ¹H NMR (DMSO-*d*₆, 270 MHz) at δ; 7.6 (2H, d, *J*= 5.7 Hz, H-2', H-6'), 7.0 (2H, d, *J*= 5.7 Hz, H-3', H-5'), 6.4 (1H, s, H-3), 6.2 (1H, d, *J*= 1.5 Hz, H-8), 6.0 (1H, d, *J*= 1.5 Hz, H-6). The EI-MS spectrum showed the molecular ion peak as the base peak

(M⁺) at *m/z* 270 (95%) corresponding to molecular formula C₁₅H₁₀O₅ together with fragments at *m/z* 152 (43%), 118 (48%), 121 (20%) and 149 (26%).

Compound (IV)

UV spectral data at λ_{max} (nm): MeOH; 265, 294, 322, 367, NaOMe; 277, 294, 315, 420, 420, AlCl₃; 268, 303, 350, 424, AlCl₃/HCl; 269, 303, 348, 424, NaOAc; 276, 302, 387, NaOAc/H₃BO₃; 266, 248, 320, 372. ¹H-NMR (DMSO-*d*₆, 270 MHz) at δ; 7.94 (2H, d, *J*= 5.4 Hz, H-2', H-6'), 6.9 (2H, d, *J*=

5.4 Hz, H-3', H-5'), 6.5 (1H, d, *J*= 1.2 Hz, H-8), 6.2 H, d, *J*= 1.2 Hz, H-6). The EI-MS spectrum showed the molecular ion peak as the base peak (M⁺) at *m/z* 287 (84%) corresponding to molecular formula C₁₅H₁₁O₆ together with fragments at *m/z* 269, 259 (12%), 258 (15%), 153 (42%) and 133 (48%)







Compound (V)

Compound is colorless crystals that gave violet color with ferric chloride solution with melting point 215-216 °. The ¹H-NMR (CD₃OD, 300 MHz) showed signals at δ; 7.58 (2H, d, *J*= 2.1 Hz, H-2, 6), 6.92 (2H, d, *J*= 1.2 Hz, H-3, 5).

Compound (VI)

Compound VI is identified by co-chromatography with authentic using paper chromatography with solvent system butanol: acetic acid: water (5:4:1) and acetic acid 6%. Spots visualized under UV light show violet color and gave blue color with FeCl₃ with R_f = 0.53 and 0.56 respectively.

Table 1: of isolated compounds

	
β-Amyrin	Lupeol
Compound (I)	Compound (II)
	
Apigenin	Kaempferol
Compound (III)	Compound (IV)
	
4-Hydroxybenzoic acid	Gallic acid
Compound (V)	Compound (VI)

4. Biological activities

4.1 Evaluation of anti-inflammatory activity

The accumulation of nitrite, an indicator of NO synthesis was measured by Griess reagent (Gerhauser, *et al.*, 2003). A standard curve was plotted using serial concentration of sodium nitrite (Fig. 1). The nitrite content was normalized to cellular protein content as measured by bicinchoninic acid assay [12, 32]. The NO inhibition percentage was calculated by submitting the nitrite content of the cell supernatant of cultures treated with DMSO (control) LPS or LPS/treated compounds (fig. 2).




Fig 1: The level of Nitric oxide in RAW 264.7 cells supernatant after the treatment with the samples (50 $\mu\text{g/ml}$) compared with LPS-treated cell (200 $\mu\text{g/ml}$), as measured by Griess assay.




Fig 2: The percentage of inhibition of Nitric oxide in LPS0stimulated RAW 264.7 cells supernatant after the treatment with the samples (50 $\mu\text{g/ml}$) compared to LPS treated cells, as measured by Griess assay.

4.2 Evaluation of antioxidant activity

DPPH is a deep violet radical due to its unpaired electron, which in the presence of antioxidant radical scavenger (donate an electron to DPPH) decolorizes to the pale yellow non-radical form [30]. The change in the colorization and subsequent fall in absorbance were monitored spectrophotometrically at

520 nm. A standard calibration curve was plotted using serial dilutions of ascorbic acid in concentrations ranging from 0 to 100 $\mu\text{g/ml}$ in distilled water (fig. 3). The half maximal scavenging capacity (SC_{50}) values for each tested sample and ascorbic acid were estimated via dose curve and calculated from curve equation (fig. 4).




Fig 3: Antioxidant activity against DPPH radicals




Fig 4: Antioxidant activity against DPPH radicals

4.3 Evaluation of cytotoxic activity

Cytotoxicity of the tested extract was measured against Hep-G₂ cells using MTT cell viability assay, which is based on ability of active mitochondrial dehydrogenase enzymes of living cells to cleave the tetrazolium rings of the yellow MTT and form dark blue insoluble formazan crystals which are largely impermeable to cell membranes, results in its accumulation within healthy cells (result in the liberation of crystals, which are then solubilized). The number of viable cells is directly proportional to the level formazan dark blue

color. The extent of reduction of MTT was quantified by measuring the absorbance at 570 nm^[14]. The percentage of relative viability was calculated using the following equation (absorbance of treated cells / absorbance of control cell)*100. Then the half maximal inhibitory concentration (IC₅₀) was calculated from the equation of dose response curve. Results showed that butanol extracts had weak cytotoxicity activity against Hep-G₂ cells, IC₅₀ = 568.5 µg/ml (fig. 5), (Olajire and azeez, 2011).




Fig 5: Cytotoxic effect of different samples against Hep-G2 cells using MTT assay (n=4), data expressed as the mean value of cell viability (% of control) \pm S.E.

5. References

- Jamal K, Yaacob WA, Laily DB. *Journal of Physical Science* 2008; 19(2):45–50.
- Qiao A, Wang Y, Xiang L, Zhang Z, Xiangjiu H. *Fitoterapia* 2014; 98:137–142.
- Alexsandro B, Angelo PC, Raimundo BF. *Anais Da Academia Brasileira De Ciências* 2004; 76(3):505-518.
- Joshi AB, Desai RR, Bhohe MP. *Der Pharma Chemica* 2013; 5(3):116-121.
- Gerdin B, Srenso E. *International Journal of Microcirculation, Clinical and Experimental* 1983; 2(1):39-46.
- Byung-Sun MIN, Miyuki T, Chao-Mei MA, Norio N, Masao H. *Chem Pharm Bull* 2001; 49(5):546-550.
- Catharina FE, De-Santos TS, Marcelo DM, Vianna F, Maria Auxiliadora Kaplan C. *Molecules* 2013; 18:4247-4256.
- Dilek ERCIL, Koray Sakar M. *Turk J Chem* 2004; 28; 133-139.
- Torres-Santos EC, Lopes D, Rodrigues OR, Carauta JPP. *Phytomedicine* 2004; 11:114–120.
- Kelly EH, Anthony RT, Dennis JB. *Journal of Nutritional Biochemistry* 2002; 13(10):572–584.
- Fairouz M, Amar Z, Narimane S, Ahmed T, Salah R. *Rec Nat Prod* 2010; 4(1):91-95.
- Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JK, Annenbaum SR. *Anal Biochem* 1982; 126:131–136.
- Guang-ying Chen, Chun-yan Dai A, Tian-shan Wang A, Cai-wu Jiang A, Chang-ri Han A, Xiao-ping Songa. *ARKIVOC* 2010, 179-185.
- Hansen MB, Nielsen SE, Berg K. *J Immunol Methods* 1989; 10:119-203.
- Herbert B, Carl D. *Dudley structure elucidation of natural products by mass spectrometry*, San Francisco, Amsterdam, london: Holden-Day, 1964.
- Jiang H, Yan S, Hui L, Benshou Y, Xia M, Yong MZ *et al. Fitoterapia* 2014; 99:86–91.
- Jie Z, Jun Y, Jicheng D, Zhen L, Lihua Z, Yushu H *et al. Analytica Chimica Acta* 2005; 532:97–104.
- Jones DL. *Palms throughout the World*. Smithsonian Books, 1995.
- Jorge SAR, Ana LS, Daniela APS, Bruno GMF, Ana VS, Vanessa MIS *et al. Studies in Natural Products Chemistry* 2014; 41:33-73.
- Markham KR. *Techniques of Flavonoid Identification*. Academic Press, London, 1982.
- Li CW, Cui CB. *Molecules* 2014; 19(12):21363-21377.
- Li-Ying Liu, Hui Chen, Chao Liu, Hong-Qing Wang, Jie Kang, Yan Li *et al. Fitoterapia* 2014; 98:254–259.
- Pan MH, Lai CS, Ho CT. *Food and Function* 2010; 1(1):15-31.
- Maria A, Athinazira, Prokopios M, Serkos HA, Alexios LS. *Emmanuel Mikros. J Agric Food Chem* 2009; 57(23):11067–11074.
- Michael GA. *Journal of General Microbiology* 1983; 129:543-546.
- Olajire A, Azeez L. *Afr J Food Sci Technol* 2011; 2:022–029.
- Patlolla JM1, Rao CV. *Curr Pharm Biotechnol* 2012; 13(1):147-55.
- Qiu-Yan S, Wei-Yan Q, Zheng-Ming L, Jie Z, Jian-Jun C, Kun G. *Food Chemistry* 2011; 128:495–499.
- Ram CD, Meena R, Surya KK, Suresh A, Mohan GB. *J Nepal Chem Soc* 2009, 23.
- Ratty AK, Sunamoto J, Das NP. *Biochem Pharmacol* 1988; 37:989–995.
- Raymond ME, Xiu-Sheng M. *International Journal of Mass Spectrometry* 2004; 231:157–167.
- Smith PK, Krohn RI, Hermanson GT, Mallia AK, Gartner FH, Provenzano MD *et al. Anal Biochem* 1985; 150:76–85.
- Venkata SPC, Indra Prakash. *Journal of Chemical and Pharmaceutical Research* 2013; 5(1):261-265.
- Zhu W, Jia Q, Wang Y, Zhang Y, Xia M. *Free Radical Biology and Medicine* 2012; 52(2):314–327.

35. Wang CZ, Mehendale SR, Calway T, Yuan CS. *Am J Chin Med* 2011; 39(4):661-671.
36. Witchuda T, Orawan T. *Thai J Pharm Sci* 2007; 31:1-8.
37. Xiao ZP, Peng ZY, Peng MJ, Yan WB, Ouyang YZ, Zhu HL. *Mini Rev Med Chem* 2011; 11(2):169-177.
38. Shi Z, Wei L, Fei-fei N, Qing Z, Jing-jing R, Jia W *et al.* *Biochemical and Biophysical Research Communications* 2009; 385:551-556.