



E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2016; 5(6): 182-184
Received: 25-09-2016
Accepted: 26-10-2016

Darakhshan Masroor

Department of Pharmacognosy,
Faculty of Pharmacy and
Pharmaceutical Sciences,
University of Karachi, Karachi-
75270, Pakistan.

Rafi Akhtar Sultan

Department of Pharmacognosy,
Faculty of Pharmacy and
Pharmaceutical Sciences,
University of Karachi,
Karachi-75270, Pakistan.

Salman Ahmed

Lecturer, Department of
Pharmacognosy,
Faculty of Pharmacy and
Pharmaceutical Sciences,
University of Karachi,
Karachi-75270, Pakistan.

Muhammad Mohtasheemul Hasan

Associate Professor, Department
of Pharmacognosy, Faculty of
Pharmacy and Pharmaceutical
Sciences, University of Karachi,
Karachi-75270, Pakistan.

Correspondence**Muhammad Mohtasheemul Hasan**

Associate Professor, Department
of Pharmacognosy, Faculty of
Pharmacy and Pharmaceutical
Sciences, University of Karachi,
Karachi-75270, Pakistan.

Near infrared spectroscopy of *Cicer arietinum* L., seeds

Darakhshan Masroor, Rafi Akhtar Sultan, Salman Ahmed and Muhammad Mohtasheemul Hasan

Abstract

NIRS is an advance, fast, reliable and non-destructive technique extensively used in chemical, food, petrochemical and pharmaceutical industries for rapid and reliable standardization. NIRS of *Cicer arietinum* L. *Desi* and *Kabuli* variety seed flours was performed for qualitative evaluation of secondary metabolites. The major suggested compounds in *Cicer arietinum Desi* and *Kabuli* variety were found as adenine, biochanin-A, biochanin-B, biohannin A 7-O- β -D-glucoside, calycosin, cerebroside, choline, ciceritol, citric acid, diadzein, fructose, garbanzol, genistein, glucose, inositol, isoliquiritigenin, liquiritigenin, methyl tetracosanoate, ononin, pyrimidine, saccharose, sissotrin, stigmaterol, trifolirhizin, β -amyryn and β -sitosterol. The presence of these compounds justifies the use of *Cicer arietinum* as nutritional food and tonic.

Keywords: NIRS, *Cicer arietinum*, secondary metabolites

Introduction

The active chemical constituents of medicinal plants are responsible for various desired therapeutic effects. Several pharmaceutical companies have initiated sophisticated plant screening programs to find out promising active therapeutic compound(s) in plants. The infrared (IR) region in spectroscopic techniques is used in the agriculture, chemical, cosmetic, food, medicine, petrochemical, pharmaceutical and textile industries. In 1800, Frederick William Herschel a British astronomer discovered the light in the near infrared (NIR) region, but the first practical application of NIRS as an analytical method was carried out by Karl Norris (an engineer working at the USDA Instrumentation Research Laboratory) on grains and seeds in 1960s. NIRS is now a recognized analytical method for grains and seeds by the American Association of Cereal Chemists and the American Oil Chemist Association. In the near infrared (NIR) region, the spectral signatures appear due to the stretching, bending and rotating vibrations in the chemical compounds^[1, 2]. NIRS is a rapid, reliable and non-destructive technique which provides acceptable accuracy with a minimum sample preparation in both qualitative and quantitative analysis to meet the requirements of quality control. This technique is successfully used to determine ash, carbohydrates, fatty acids, inorganic phosphorus, moisture, protein, starch and total dietary fiber^[1, 3-5].

Cicer arietinum belongs to family Fabaceae, commonly known as Chickpea in English and *Channa* in Urdu. Chickpea is distributed in Africa, Asia, Caribbean, Central America, Europe, Middle East and South America. *Desi* (black) and *Kabuli* (white) *Channa* are two cultivated varieties based on color and size. The chickpea seeds have carbohydrates, proteins (alanine, arginine, aspartic acid, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tyrosine and valine amino acids), fatty acids (arachidic, linoleic, myristic, oleic, palmitic and stearic acids) and vitamins A-E. In secondary metabolites, saponins, phenolic acids (caffeic acid, chlorogenic acid, ferulic acid, vanillic acid) and β -sitosterol are reported^[6, 7]. The seeds have also shown antibacterial, anti-carcinogenic, antidiabetic, antifungal, anti-inflammatory, antioxidant, diuretic, hepatoprotective, hypocholesterolemic, hypoglycemic and purgative activities^[6-9].

Materials and methods**Plant material, identification and sample preparation**

The seeds of *Cicer arietinum* L., were purchased and authenticated by a taxonomist of the Department of Botany, University of Karachi. The voucher specimen number (CAB-06-15/16) and (CAW-07-15/16) were deposited in the Herbarium. The seeds were separately grinded and powdered then passed through 600 μ m sieve and kept in an amber bottle at room temperature before experiment.

NIR spectral data

FOSS XDS near infrared Rapid Content Analyzer (FOSS NIR Systems, Inc., USA) was used for NIR spectrum (700 - 2500 nm). XDS RCM Iris was used to center the sample for reproducible sample positioning. NIR reflectance spectra were recorded on FOSS XDS near infrared Rapid Content Analyzer at ambient temperature. Powdered sample (10g) was filled in a disposable glass silicate vial. Prior to the screening, the vial was closed with a plastic lid, then inverted once and tapped twice on a soft surface. The outer surface of the glass vial was clean from any contamination before analysis. The vial then placed onto the sample presentation glass of Foss Rapid Content Analyzer and centered by using the Iris centering device. The Iris was retracted completely after centering and before scanning [10].

Results and Discussion

The NIR spectrum of *C. arietinum Desi* and *Kabuli* variety suggested the presence of adenine, β -amyrin, β -sitosterol, biochanin A 7-O- β -D-glucoside, biochanin A, biochanin B, calycosin, cerebroside, choline, ciceritol, citric acid, diadzein, fructose, garbanzol, genistein, glucose, inositol, isoliquiritigen, liquiritigenin, methyl tetracosanoate, ononin, pyrimidine, saccharose, sissotrin, stigmaterol and trifolirhizin. The results of NIRS are presented in table 1-2 and figure 1-2.

NIR spectroscopy is useful analytical technique to assess the

structure of organic compounds containing C-H bonds, O-H, and N-H bonds through the absorption of energy in the NIR region of the spectrum due to relatively weak, broad overtones and combination bands of fundamental vibrations transitions. It has been effectively used for the analysis of natural products and for assessment of product quality [1, 3]. In this study, the spectral data are analyzed between the regions of 1100 – 2400 nm. Each and every peak's wave length of specific spectrum indicates possible functional groups, thus provide fingerprint of a compound in taking powdered plant material. The detail of NIR spectra with specific wavelengths, bond vibrations, possible functional groups and the compounds responsible for the peak present in the seed flour of *C. arietinum* are given in table. The data obtained confirms the presence of chemical compounds reported in the literature.

Conclusion

The results of NIR analysis of the powder confirms the presence of reported chemical compounds. The compounds justify the nutritional importance of *C. arietinum*. This technique can be used as a rapid and authentic method for herbal drug standardization.

Conflict of Interest

The authors declare no conflict of interest regarding the publication of this paper.

Table 1: Wave length of spectral peaks obtained from *Cicer arietinum (Desi)* seed flour and possible chemical compounds

Band vibrations with peak's wave length (nm)	Possible functional groups	Compounds present in the seed flour that may be responsible for peak
CH second overtone (1205)	CH ₂ , CH	adenine, biochanin-A, biochanin-B, biochannin a 7-o- β -d-glucoside, calycosin, cerebroside, choline, ciceritol, citric acid, diadzein, fructose, garbanzol, genistein, glucose, inositol, liquiritigenin, ononin, pyrimidine, saccharose, sissotrin, stigmaterol, trifolirhizin, β - amyryn and β -sitosterol.
First overtone of CH combinations; NH and OH first overtone (1460)	CH, CONH ₂ , ROH	
CH first overtone (1765)	CH	
N-H+C-H combination band (2125)	CC, RNH ₂	
C-H+C-H combinations band (2285, 2310, 2345)	CH ₃ , CH ₂ , CH	

Table 2: Wave length of spectral peaks obtained from *Cicer arietinum (Kabuli)* seed flour and possible chemical compounds

Band vibrations with peak's wave length (nm)	Possible functional groups	Compounds present in the seed flour that may be responsible for peak
CH second overtone (1725)	CH ₂ , CH	adenine, biochanin-A, biochanin-B, biochannin a 7-o- β -d-glucoside, calycosin, cerebroside, choline, ciceritol, citric acid, diadzein, fructose, garbanzol, genistein, glucose, inositol, liquiritigenin, ononin, pyrimidine, saccharose, sissotrin, stigmaterol, trifolirhizin, β - amyryn and β -sitosterol.
OH and NH first overtone (1470)	CH, CONH ₂ , ROH	
CH and SH first overtone (1770)	CH	
C-H+C-H combinations band (2290, 2320)	CH ₃ , CH ₂ , CH	
N-H+C-H combination band (2120)	CONH ₂ (R)	




Fig 1: NIR spectrum of *Cicer arietinum (Desi)* seed flour




Fig 2: NIR spectrum of *Cicer arietinum* (Kabuli) seed flour.

References

1. Cozzolino D. Near infrared spectroscopy in natural products analysis. *Planta Medica*, 2009; 75(07):746-756.
2. Agelet LE, Hurburgh CR. Limitations and current applications of Near Infrared Spectroscopy for single seed analysis. *Talanta*, 2014; 121:288-299.
3. Teye E, Huang X-Y, Afoakwa N. Review on the potential use of near infrared spectroscopy (NIRS) for the measurement of chemical residues in food. *American Journal of Food Science and Technology*, 2013; 1(1):1-8.
4. Plans M, Simó J, Casañas F, Sabaté J, Rodríguez-Saona L. Characterization of common beans (*Phaseolus vulgaris* L.) by infrared spectroscopy: comparison of MIR, FT-NIR and dispersive NIR using portable and benchtop instruments. *Food Research International*. 2013; 54(2):1643-1651.
5. Zhang C, Su J. Application of near infrared spectroscopy to the analysis and fast quality assessment of traditional Chinese medicinal products. *Acta Pharmaceutica Sinica B*. 2014; 4(3):182-192.
6. Bagri P, Ali M, Sultana S, Aeri V. A new flavonoid glycoside from the seeds of *Cicer arietinum* Linn. *Acta Poloniae Pharmaceutica*, 2011; 68(4):605-608.
7. Al-Snafi AE. The medical Importance of *Cicer arietinum* -A review. *IOSR Journal of Pharmcay*, 2016; 6(3):29-40.
8. Jukanti AK, Gaur PM, Gowda CLL, Chibbar RN. Nutritional quality and health benefits of chickpea (*Cicer arietinum* L.): a review. *British Journal of Nutrition*, 2012; 108:S11-S26.
9. Tiwari AK, Sahana C, Zehra A, Madhusudana K, Kumar DA, Agawane SB. Mitigation of starch-induced postprandial glycemic spikes in rats by antioxidants-rich extract of *Cicer arietinum* Linn. seeds and sprouts. *Journal of Pharmacy & Bioallied Sciences*. 2013; 5(4):270-276.
10. Foss NIR Systems. Installation and user guide, Foss Rapid Content Analyzer TM, in Part Number 720-750-6540; Version 1.0. U.S.A.: Foss NIR Systems, Inc. 1998.