



E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2017; 6(6): 738-741
Received: 18-09-2017
Accepted: 19-10-2017

Avadh Bihari Singh
Department of Vegetable
Science, B. C. K. V. Mohanpur,
Nadia, West Bengal, India

Chandra Deo
Department of Vegetable
Science, B. C. K. V. Mohanpur,
Nadia, West Bengal, India

Sanket Kumar
Department of Vegetable
Science, B. C. K. V. Mohanpur,
Nadia, West Bengal, India

Shiv Prakash Shrivastav
Genetics and Plant Breeding,
NDUA & T, Kumarganj,
Faizabad, Uttar Pradesh, India

Sriom
Department of Vegetable
Science, B. C. K. V. Mohanpur,
Nadia, West Bengal, India

Correspondence
Avadh Bihari Singh
Department of Vegetable
Science, B. C. K. V. Mohanpur,
Nadia, West Bengal, India

Growth and yield response of sweet potato (*Ipomoea batatas* L.) cv. NDSP-65 to different integrated organic sources

Avadh Bihari Singh, Chandra Deo, Sanket Kumar, Shiv Prakash Shrivastav and Sriom

Abstract

A field experiment was conducted in the *rabi* season at Department of Vegetable Science, N.D.U.A. & T., Faizabad (U.P.) during 2015-16, to find out the suitable doses of organic manures for sustainable production of Sweet Potato (*Ipomoea batatas* L.). The total of eleven doses of organic manures along with different combinations was evaluated by using randomized block design. Earliest initiation of buds (7.93 days) was observed in T₁ (FYM @ 20 t/ha). T₁₁ {recommended dose of FYM (10 t/ha) and NPK (50:25:50)} was identified as a good combination for number of leaves per plant, leaf area, foliage weight per plant, number of vine per plant, inter nodal length, number of tuber plant, fresh weight of tuber per plant, length of tubers and yield per hectare. However for all organic sources, treatment T₇ {FYM @ 10 t / ha + Poultry manure @ 2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} could be used for improving growth and yield along with related traits of sweet potato. Hence, it is suggested that these remunerative treatment of organic doses help in successful crop production of sweet potato.

Keywords: sweet potato (*Ipomoea batatas* L.), NDSP-65, organic manures, growth, tuber yield

Introduction

Organic manures are the most important input that significantly increases the productivity of the crop (Panwar and Wani, 2014) [25]. It can also improve the nutrients and is absolutely harmless to plants. Sweet potato [*Ipomoea batatas* (L.) Lam] is important root crops, belongs to family convolvulaceae, considered to be widely grown in tropical and sub-tropical regions of the world (Onunka *et al.*, 2012) [22]. In India, sweet potato area, production and productivity were 0.11 million hectares, 1.45 million tonnes and 13.06 tonnes per hectare, respectively (Anonymous, 2015) [6]. The sweet starchy edible tuberous roots have economic values that contain about 27% carbohydrate and high concentration of Vitamin A, Vitamin C, calcium and iron. Sometimes, young leaves and shoots are also eaten as greens (Abidin, 2004) [2]. It is a rich source of carotene, ascorbic acid, thiamine, riboflavin, protein and energy. Because of the high nutritional and economic value, it is necessary to improve yield and its related traits that can be achieved through balance availability of all the nutrients in the crop. No single source is capable to supply the required quantity of plant nutrients. Due to continuous application of inorganic fertilizers in the soil, had negative effect on soil and also productivity of the crop (Lal and Kang, 1982) [14].

There is large number of organic sources such as farm yard manure, town compost, horse manure, sewage sludge, press mud, goat and sheep manure, cattle manure, vermi-compost etc. These could be used to reduce the total cost of cultivation and to supplement the essential nutrients for better growth and development of the plants. In recent past research, it is found that the root yield of sweet potato increases with application of inorganic fertilizers but adversely affects the quality (Nedunchezhiyan and Srinivasulu Reddy, 2002) [17]. However, optimum amount of nitrogen particularly organically increases quality (Nedunchezhiyan *et al.*, 2003) [18].

Bio-fertilizers is one of the most important organic source, contain beneficial viable-organisms which have ability to mobilize nutritionally important elements from non-usable to usable form through biological processes (Oliveira *et al.*, 2010) [21]. *Azospirillum* is known to be a very active nitrogen fixer under laboratory as well as soil condition providing fast growth, better health of the plant and higher yield (Kannan and Ponmurugan, 2010) [12]. The response of organic sources with or without chemical fertilizers on a large number of crops have been reported by several workers (Nedunchezhiyan *et al.*, 2010; Sowley *et al.*, 2015; Koodi *et al.*, 2017) [19, 29, 13].

Considering this, the present study was undertaken to find out the suitable organic sources for higher yield of sweet potato with the economic feasibility of the treatments.

Materials and Methods

The experiment was carried out during 2015-2016, at Main Experiment Station, Department of Vegetable Science, N.D.U.A. & T., Faizabad (U.P) India. The experimental site falls under sub-humid, subtropical climate and is located at 26.47° N latitude and 82.12° E longitudes on an elevation of 113 meters above mean sea level in the Indo-gangetic alluvial plains of eastern Uttar Pradesh. Maximum rainfall in this area is received from mid-June to end of September. The weekly maximum and minimum temperatures during the crop growth period ranged from 36.6 and 20.1 and 25.8 to 5.2, respectively. The total rainfall recorded during the crop period was 15.2 mm.

A well prepared manured nursery having good drainage helps in producing better planting materials. Planting in nursery is done 3 months before ahead of planting in main field. The selected tubers are planted 5-10 cm deep at a spacing of 30 cm in rows, 60 cm a part. The sprouts are often cut after 40-45 days and planted in secondary planted nursery for further growth. The vine cutting, 20-30 cm in length, the cuttings are made and planted in the field, with a spacing of 60 cm x 30 cm. All the recommended agronomic package of practices and plant protection measures were followed to raise a good crop.

The crop was harvested on 4th and 5th February during the year 2015-2016.

The recommended dose of NPK was 50:25:50 kg/ha and 25:12.5:25 kg/ha. Nitrogen fertilizers were applied as per treatments under study at the last ploughing, the whole quantity of organic manure, vermicompost and biofertilizer was incorporated in the soil as per treatment under study. The different treatments are given in table 1 with the details of doses.

Observations recorded on five randomly selected plants from each genotype in each replication for growth and yield along with related characters viz., days to initiation of buds, number of leaves per plant, leaf area, foliage weight per plant, number of vine per plant, inter nodal length, tuber weight, number of tubers per plant, fresh weight of tuber per plant, length of tubers, diameter of tubers and yield per hectare. The collected data were averaged to get mean values of the respective characters that has been affected by various treatments integrated nutrient managements in sweet potato.

The data were subjected to the analysis of variance (ANOVA) appropriate the design and test of significance of the treatment difference was done on the basis of F test (Gomez and Gomez, 1984). The treatments were compared with the help of critical difference, following the techniques described by (Panse and Sukhatme, 1967) [24] and results were evaluated at 5% level of significance.

Table 1: Different integrated organic treatments with their respective doses

Treatment	Doses
T ₁	FYM @ 20 t/ha
T ₂	Poultry manure @ 5t/ha
T ₃	Neem cake @ 4 t/ha + <i>Azospirillum</i> (5 kg/ha) + PSB (5 kg/ha)
T ₄	Vermicompost @5 t/ha + <i>Azospirillum</i> (5 kg/ha) + PSB (5 kg/ha)
T ₅	FYM @ 10 t/ha + Vermicompost 2.5 t/ha + <i>Azospirillum</i> (5 kg/ha) + PSB (5 kg/ha)
T ₆	FYM @ 10 t/ha + Neem cake @1 t/ha + <i>Azospirillum</i> (5 kg/ha) + PSB (5 kg/ha)
T ₇	FYM @ 10 t/ha + Poultry manure @2.5 t/ha + <i>Azospirillum</i> (5 kg/ha) + PSB (5 kg/ha)
T ₈	Vermicompost @ 2.5 t/ha + Neem cake @ 1 t/ha + <i>Azospirillum</i> (5 kg/ha) + PSB (5 kg/ha)
T ₉	Vermicompost @ 2.5 t/ha + Poultry manure 2.5 t/ha + <i>Azospirillum</i> (5 kg/ha) + PSB (5 kg/ha)
T ₁₀	1/2 Recommended dose of Fertilizers + <i>Azospirillum</i> (2.5 kg/ha) + PSB (2.5 kg/ha)
T ₁₁	Recommended dose of FYM and NPK (10 t/ha and 50:25:50 kg NPK/ha)

Results and discussion

Growth parameters

The effect of different treatments on the growth parameters of sweet potato are given in Table 2. Application of different organic and inorganic fertilizers rates leads to significant differences in most the growth parameters except days taken to bud initiation.

It was observed that the days taken to bud initiation influenced non-significantly due to various organic treatments. However, a higher day to initiation of buds was recorded under T₁ (FYM @ 20 t/ha) followed by T₂ (Poultry manure @ 5t/ha) and T₄ {Vermicompost @5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} as compared rest of the organic treatments. Data also revealed that T₇ {FYM @ 10 t/ha + Poultry manure @2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} treatments recorded lowest number of days taken to initiation of buds. These results are in accordance with Abdissa *et al.*, (2012) [1].

Treatment T₁₁ {Recommended dose of FYM and NPK (10 t/ha and 50:25:50 kg NPK/ha)} recorded maximum number of leaves was statistically at par with T₇ {FYM @ 10 t/ha + Poultry manure @2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} treatment and significantly better over rest of the treatments. The minimum number of leaves was recorded in

treatment T₂ (Poultry manure @ 5t/ha) with a value of 180.60. Mukhtar *et al.*, (2010) [15] and Adeyeye *et al.*, (2016) [3] also reported the similar findings.

The maximum size of leaf area of (30.30 cm²) was found in treatment T₁₁ {Recommended dose of FYM and NPK (10 t/ha and 50:25:50 kg NPK/ha)} which was statistically at par with treatments T₇ {FYM @ 10 t/ha + Poultry manure @2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} and T₆ {FYM @ 10 t/ha + Neem cake @1 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} and significantly superior over rest of the treatments. These kind of results also obtained by Ghosh and Das (1998) [9]; Sood and Sharma (2001) [28]; Amara and Mourad (2013) [5]; Koodi *et al.*, (2017) [13].

The treatment T₁₁ {Recommended dose of FYM and NPK (10 t/ha and 50:25:50 kg NPK/ha)} produced maximum foliage weight per plant, number of vine per plant and inter nodal length with mean value of 432.90 g, 10.68 and 4.10 cm respectively, which was statistically at par with T₇ {FYM @ 10 t/ha + Poultry manure @2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} and T₆ {FYM @ 10 t/ha + Neem cake @1 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)}. Higher levels of nutrients and a helped in cell elongation of stem due to development of cell and rapid cell division and cell elongation in meristematic region of plant. Similar results were also

reported by Agbede (2011) [4] Nongmaithem and Pal (2011) [20].

The minimum number of vine per plant was recorded in T₄ {Vermicompost @5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)}. However, the minimum foliage weight per plant and

inter nodal length were found with the application of T₂ (Poultry manure @ 5t/ha). Similar findings has been reported by Panigraphi and Behera (1993) [23]; El Gamal (1996) [8]; Sood and Sharma (2001) [28]; Panwar and Wani (2014) [25].

Table 2: Effect of different integrated organic treatments for growth, yield and its related traits of sweet potato

Treatment	Days to initiation of buds	Number of leaves per plant	Leaf area (cm ²)	Foliage weight per plant	Number of vine per plant	Inter nodal length (cm)	Number of tuber per plant	Tuber Weight (g)	Fresh weight of tuber per plant (g)	Length of tubers(cm)	Diameter of tubers (cm)	Yield per hectare (q)
T ₁	7.93	205.80	25.38	362.60	8.13	3.43	3.50	98.00	343.00	13.52	5.15	201.39
T ₂	7.95	180.60	22.27	318.20	6.52	3.01	3.00	100.33	301.00	11.87	6.35	176.73
T ₃	8.20	201.60	24.86	355.20	7.86	3.36	3.40	98.82	336.00	13.25	5.17	197.28
T ₄	8.03	186.90	23.05	329.30	5.73	3.14	3.15	98.89	311.50	12.28	5.31	182.90
T ₅	9.00	220.60	27.45	392.20	9.20	3.71	3.40	109.12	371.00	14.63	6.47	217.83
T ₆	9.27	228.90	28.23	403.30	9.61	3.83	4.00	95.38	381.50	15.24	4.97	224.00
T ₇	9.50	235.20	29.01	414.40	10.01	3.92	3.60	108.89	392.00	15.46	6.1	230.16
T ₈	8.60	197.40	24.35	347.80	7.60	3.29	3.20	102.81	329.00	12.97	5.11	193.17
T ₉	8.33	191.10	23.57	336.70	7.19	3.19	3.00	106.17	318.50	12.56	6.3	187.01
T ₁₀	8.50	214.20	26.42	377.40	8.67	3.57	3.50	102.00	357.00	14.08	5.7	209.68

Yield parameters

Yield per hectare and their related characters were affected by various integrated organic treatments have been presented in Table 2. Different treatments have their significant impact of number of tubers per plant. The maximum number of tubers (4.0/plant) were noted under 2 treatments viz., T₁₁ {Recommended dose of FYM and NPK (10 t/ha and 50:25:50 kg NPK/ha)} and T₆ {FYM @ 10 t/ha + Neem cake @ 1 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha) + PSB (5 kg/ha)} treatment which was at par with T₇ {FYM @ 10 t/ha + Poultry manure @ 2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)}. It is obvious from the table that minimum value was noted under T₂ (Poultry manure @ 5t/ha). These kind of results also obtained by Imam and Badawy (1978) [11]; Yadav *et al.*, (2003) [30]; Raghav and Kamal (2009) [26]; Sharma and Sharma (2011) [27].

The perusal of data obviously indicated that tuber weight influenced significantly due to various organic treatments. T₅ {FYM @ 10 t/ha + Vermicompost 2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} treatment produced maximum weight of tuber (109.12 g) followed by T₇ {FYM @ 10 t/ha + Poultry manure @ 2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} and T₈ {Vermicompost @ 2.5 t/ha + Neem cake @ 1 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)}. However, minimum tuber weight *i.e.* 95.38 g was observed in the treatment T₆ {FYM @ 10 t/ha + Neem cake @ 1 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha) + PSB (5 kg/ha)}. Raghav and Kamal (2009) [26]; Zaman (2011) [32] also reported the similar findings.

Treatment T₁₁ {recommended dose of FYM and NPK (10 t/ha and 50:25:50 kg NPK/ha)} produced maximum fresh weight of tuber per plant (409.50 g) that was statistically at par with T₇ {FYM @ 10 t/ha + Poultry manure @ 2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} and significantly superior over the rest of treatments. The minimum fresh tuber weight per plant was observed with the treatment T₂ (Poultry manure @ 5t/ha). These results are in accordance with Mukhtar *et al.*, (2010) [15]; Yourtchi *et al.*, (2013) [31]; Adeyeye *et al.*, (2016) [3].

The maximum length of tuber (16.15) cm was noted with treatment T₁₁ {Recommended dose of FYM and NPK (10 t/ha and 50:25:50 kg NPK/ha)} which was statistically at par with T₇, and T₆ and significantly superior over rest of the treatments. However, the minimum length of tuber *i.e.* 11.87

cm was observed in plots receiving T₂ (Poultry manure @ 5t/ha). Treatments T₅ {FYM @ 10 t/ha + Vermicompost 2.5 t/ha + *Azospirillum* 5 kg/ha + PSB 5 kg/ha} produced maximum diameter of tubers (6.47) cm which was significantly superior over the rest of treatments. Similar findings has been reported by Yourtchi *et al.*, (2013) [31].

Highest tuber yield per ha (240.44 q/ha) was recorded under treatment T₁₁ {recommended dose of FYM and NPK 10 t/ha and 50:25:50 kg NPK/ha} which was significantly at par with T₇ {FYM @ 10 t / ha + Poultry manure @ 2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} and T₅ {(FYM @ 10 t / ha + RDF + 1 t/ha Neem cake)}. Similar results were also reported by Byju and Ravindran (2009) [7]; Yadav *et al.*, (2003) [30]; Raghav and Kamal (2009) [26]; Zaman (2011) [32]; Narayan *et al.*, (2013) [16]; Nedunchezhiyan *et al.*, (2010) [19]; Sowley *et al.*, (2015) [29]. However, minimum tuber yield per hectare *i.e.* 176.73q was recorded under T₂ {Poultry manure @ 5t/ha} treatment.

Conclusion

Therefore, it is concluded that, T₁₁ {Recommended dose of FYM and NPK (10 t/ha and 50:25:50 kg NPK/ha)} was observed best treatment for growth related traits. For yield related traits, treatment T₁₁ {Recommended dose of FYM and NPK (10 t/ha and 50:25:50 kg NPK/ha)} could be more significant for number of tuber per plant, fresh weight of tubers per plant, length of tubers and yield per hectare. Whereas T₅ {FYM @ 10 t/ha + Vermicompost 2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} for tuber weight and diameter of tubers. While in case of all organic sources, treatment T₇ {FYM @ 10 t / ha + Poultry manure @ 2.5 t/ha + *Azospirillum* (5 kg/ha) + PSB (5 kg/ha)} might be used for improving growth and yield along with related traits of sweet potato.

References

1. Abdissa T, Dechassa N, Alemayehu Y. Sweet Potato Growth Parameters as Affected by Farmyard Manure and Phosphorus Application at Adami Tulu, Central Rift Valley of Ethiopia. Agricultural Science Research Journal. 2012; 2(1):1-12.
2. Abidin PE. Sweet potato breeding for northeastern Uganda: Farmer varieties, farmer-participatory selection, and stability of performance (PhD Thesis). The

- Netherlands: Wageningen University. 2004, 152.
3. Adeyeye AS, Akanbi WB, Sobola OO, Lamidi WA, Olalekan KK. Comparative Effect of Organic and In-Organic Fertilizer Treatment on the Growth and Tuber yield of Sweet Potato (*Ipomea Batata* L.). International Journal of Sustainable Agricultural Research. 2016; 3(3): 54-57.
 4. Agbede TM, Adekiya AO. Evaluation of sweet potato (*Ipomoea batatas* L.) performance and soil properties under tillage methods and poultry manure levels. Emir. J Food Agric. 2011; 23(2):164-177.
 5. Amara DG, Mourad SM. Influence of organic manure on the vegetative growth and tuber production of potato (*solanumtuberosum* L varspunta) in a Sahara desert region. International Journal of Agriculture and Crop Sciences, 2013. 013/5-22: 2724-2731.
 6. Anonymous. Horticulture Data Base, National Horticulture Board, Ministry of Agriculture and Farmers's Welfare. Gurgaon, Haryana, 2015.
 7. Byju G, Ravindran CS. Effect of *Azospirillum* in increasing yield and nitrogen use efficiency of sweet potato cultivation in India. Advances in Horticultural Science. 2009; 23(4):254-258
 8. Gamal El, AM. Response of potato to phosphorus fertilizer levels and phosphorinebiofertilizer in the newly reclaimed areas. Assiut. J Agri. Sci. 1996; 27(2):77-87.
 9. Ghosh DC, Das AK. Effect of biofertilizers and growth regulators on growth and productivity of potato (*Solanumtuberosum*L.) Ind. Agri. 1998; 42(2):109-113.
 10. Gomez KA, Gomez AAS. Statistical procedures for agricultural research. John Wiley & Sons, New York, 1984.
 11. Imam MK, Badawy FH. Response of three potato cultivars to inoculation with *Azotobacter*. Potato Research. 1978; 21(1):1-8.
 12. Kannan T, Ponmurugan P. Response of paddy (*Oryza sativa* L.) varieties to *Azospirillum brasilense* inoculation. Journal of Phytoogy. 2010; 2(6):08-13.
 13. Koodi S, Singh SP, Rolaniya MK, Raj P. The Growth, yield and quality of sweet potato (*Ipomoea batatas* Lam.) Influenced by different plant densities International Journal of Chemical Studies. 2017; 5(4):359-361.
 14. Lal R, Kang BT. Management of organic matter in soils of the tropics and subtropics. Pages 152-178 in Non-symbiotic nitrogen fixation and organic matter in the tropics. Symposia papers, 12th congress of the International Soil Science Society, New Delhi, India, 1982.
 15. Mukhtar AA, Tanimu B, Arunah UL, Babaji BA. Evaluation of the Agronomic Characters of Sweet Potato Varieties Grown at Varying Levels of Organic and Inorganic Fertilizer. World Journal of Agricultural Sciences. 2010; 6(4):370-373.
 16. Narayan S, Kanth RH, Khan FA, Narayan R., Nabi, A. Mir SA. Growth and yield response of potato to date of planting and integrated nutrient management. Ind. J. Plt. Physio. 2013; 18 (3):270-276.
 17. Nedunchezhiyan M, Srinivasulu Reddy D. Nitrogen management in sweet potato (*Ipomoea batatas* L.) under rainfed conditions. Indian J. Agron. 2002; 47(3):449-454.
 18. Nedunchezhiyan M, Srinivasulu Reddy D, Haribabu K. Nitrogen management practices on quality characters of sweet potato (*Ipomoea batatas* L. Lam). J. Root Crops. 2003; 29(2):69-72.
 19. Nedunchezhiyan M, Byju G, Dash SN. Effects of organic production of orange fleshed sweet potato (*Ipomoea batatas* L.) on root yield, quality and soil biological health. International Research Journal of Plant Science. 2010; 1(6):136-143.
 20. Nongmaithem D, Pal D. The effect of organic sources of nutrients on the growth attributes and yields of potato (*Solanumtuberosum* L.). J Crop and Weed. 2011; 7(2):67-69.
 21. Oliveira AP, Santos JF, Cavalcante LF, Pereira WE, Santos MC, Oliveira ANP. *Et al.* Yield of sweet potato fertilized with cattle manure and biofertilizer Horticultura Brasileira. 2010; 28(3).
 22. Onunka NA, Chukwu LI, Mbanasor EO, Ebeniro CN. Effect of organic and inorganic manures and time of application on soil properties and yield of sweet potato in a tropical ultisol. Journal of Agriculture and Social Research. 2012; 12(1):183-193.
 23. Panigrahi UC, Behera B. Response of *Azotobacter*inoculants on total nitrogen, organic carbon, and microbial population of soil and yield of potato. Ind. J Agri. Chem. 1993; 26(1):17-23.
 24. Panse VG, Shukhatme PV. Statistical methods for agriculture workers (2nd eds.). Indian Council of Agriculture Research, New Delhi, 1967.
 25. Panwar S, Wani AM. Effect of Organic production on growth and productivity of Sweet Potato (*Ipomoea batatas* L.) under Poplar based Agroforestry system. International Journal of Advanced Research. 2014; 2(12):229-232.
 26. Raghav M, Kamal S. Effect of organic sources of nutrients on potato production in Tarai region of Uttarakhand. Pantnagar J Res. 2009; 7(1):69-72.
 27. Sharma JS, Sharma AK. Effect of organic fertilizers on growth, yield and quality of potato under rainfed conditions of central Himalayan region. Potato J. 2011; 38(2):176-181.
 28. Sood MC, Sharma RC. Value of growth promoting bacteria, vermicompost and *Azotobacter* on potato production in Shimla hills. J Ind. Potato Assoc. 2001; 28(1):52-53.
 29. Sowley ENK, Neindow M, Abubakari AH. Effect of poultry manure and NPK on yield and storability of orange-and white-fleshed sweet potato [*Ipomoea batatas* (L.) Lam]. Journal of Food and Agriculture Science. 2015; 5(1):1-6.
 30. Yadav KS, Singh N, Suneja S, Malik YS, Nehra B. Narula N. Nitrogen economy through the use of *Azotobacter* biofertilizer in potato. J Ind. Potato Assoc. 2003; 30(1/2):81-82.
 31. Yourtchi MS, Hadi MHS, Darzi MT. Effect of nitrogen fertilizer and vermicompost on vegetative growth, yield and NPK uptake by tuber of potato (cv. Agria). Int. J Agri. and Crop Sci. 2013; 5(18):2033-2040.
 32. Zaman A, Sarkar A, Sarkar S, Devi WP. Effect of organic and inorganic sources of nutrients on productivity, specific gravity and processing quality of potato (*Solanumtuberosum* L.). Ind. J Agri. Sci. 2011; 81(12):1137-1142.