



E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2019; SP1: 288-290

Sukhjinder Singh
Department of floriculture and
vegetable science, Baba Farid
College, Bathinda, Punjab, India

Amit Saurabh
Assistant professor Department
of Agriculture Baba Farid
College Baba Farid group of
institutions Bathinda, Punjab,
India

Ashutosh Sharma
Assistant professor Department
of Agriculture Baba Farid
College Baba Farid group of
institutions Bathinda, Punjab,
India

Correspondence
Sukhjinder Singh
Department of floriculture and
vegetable science, Baba Farid
College, Bathinda, Punjab, India

(Special Issue- 1)
2nd International Conference

“Food Security, Nutrition and Sustainable Agriculture -
Emerging Technologies”
(February 14-16, 2019)

Performance of onion under pre & post-emergence Application of herbicides and hand weeding for weed management

Sukhjinder Singh, Amit Saurabh and Ashutosh Sharma

Abstract

The present study entitled “Performance of onion (*Allium cepa* L.)” variety (N-53) under pre-emergence and post-emergence application of herbicides for weed management was carried out at CSKHPKV, Hill Agri. Research and Extension centre, Dhaulakuan (Sirmour), during *Rabi* season 2013-14 with the objectives to find out the efficacy of Pre & Post treatments on the growth, yield and yield attributing characters. In this study, 9 treatments *viz.*, Oxyfluorfen @ (0.10 kg ha⁻¹) *fb* one hand weeding (60 DAT), Oxyfluorfen @ 0.20 kg ha⁻¹, Oxyfluorfen @ 0.30 kg ha⁻¹, Pendimethalin @ 1.5 kg ha⁻¹, two hand weeding (40 and 60 DAT), farmer practice (three hand weeding at 30, 60, 90 DAT), weed free and weedy check were compared in a Randomized Block Design (RBD) having three replications with a plot size of 3.0 m x 2.0 m and a plant spacing of 20 cm x 10 cm. All the treatments were found significant for different characters. Maximum value for plant height, average weight of bulb (80 g), diameter of bulb (6.34 cm), plant population (276.33) and bulb yield (272.22 q ha⁻¹) were recorded with weed free treatment followed by Oxyfluorfen @ 0.30 kg ha⁻¹ among all herbicidal treatments. Hence, among herbicidal application Oxyfluorfen @ 0.30 kg ha⁻¹ was found most effective for most of the character studied.

Keywords: Performance, onion, herbicides, herbicides

1. Introduction

Onion (*Allium cepa* L.) is one of the most important vegetable crops in all over the world including India. It is most widely grown and popular vegetable crop among the *alliums*, globally it is considered to be the second most important vegetable after tomatoes. It is an indispensable item in every kitchen as vegetable. It is consumed as fresh salad and / or added as a spice while cooking dishes. Apart from furnishing nutrition, it also provides relishing flavors to our diets. Therefore, onion is popularly referred as ‘Queen of the kitchen’. Onion contains carbohydrates (11.0g), proteins (1.2g), fiber (0.6 g), moisture (86.8 g) and several vitamin like vitamin A (0.012 mg), vitamin C (11 mg), thiamin (0.08 mg), riboflavin (0.01 mg) and niacin (0.2 mg) and also some minerals like phosphorus (39 mg), calcium (27 mg), sodium (1.0 mg), iron (0.7 mg) and potassium (157 mg) (Rahman *et al.*, 2012)^[9].

According to Vavilov (1951)^[13], the primary center of origin lies in Central Asia. The near east and Mediterranean is the secondary centre of origin. Globally, onion is grown in an area of 44.43 lakh hectares with annual production of 856.10 lakh tonnes. (Anonymous, 2014)^[1]. India is the second largest producer and third largest exporter of onion in the world. Maharashtra is the leading producer of onion in India. In India onion is grown over an area of 12.03 lakh hectares with a production of 194.01 lakh tonnes and in Himachal Pradesh onion is grown over an area of 2340 hectares with a production of 43710 metric tonnes (Anonymous, 2014)^[1].

Weeds poses serious production problems in onion fields. The weeds problem is becoming worse in irrigated areas where cropping intensity is rapidly increasing and weed management through cultivation practices has become a challenge. They compete with onion for light, nutrient, water space and also host plant of several harmful insects and pathogens and

considerably reduce the yield, quality and value of the crop through increased production and harvesting cost (Uygun *et al.* 2010) [12]. Weed interferes with the development of onion bulbs thereby reducing bulbs yield to the extent of 40-80% (Singh *et al.* 1992) [11].

Weeds can be controlled by cultural, mechanical, chemical methods and the use of organic and inorganic mulch. On the other hand, herbicides are important tool for weed control, but are not effective in controlling the entire weed present in the crop. Use of herbicide and cultural practices as an integrated approach may provide economically acceptable weed control.

2. Material and Methods

The present investigation on "Integrated weed management studies in onion (*Allium cepa* L.)" was conducted at the Regional Research Station, Dhaulakuan district Sirmour during Rabi 2013-14. The experimental site is located at

30.40 N latitude and 77.50 E longitude at an elevation of 464 meters above mean sea level. The climate of Dhaulakuan, Himachal Pradesh is generally sub-temperate to sub-tropical; May and June are the hottest months and December and January are the coldest ones. In the year 2014, during the cropping season i.e. January to May, maximum mean weekly temperature ranged from 17.6 to 40.9 °C and minimum from 4.9 to 19.1 °C while relative humidity varied from 51.9 to 85.4 per cent. The crop experienced well distributed rainfall of 182.2 mm during the crop season. The highest weekly rainfall of 33.2 mm was received during meteorological week 9. The sunshine duration ranged from 1.5 to 10.8 hrs. The present investigation was conducted in randomized block design with three replications. Including weedy check and weed free there were in all 9 treatments. The details of treatments have been given here as below:

Table 1

Sr. No	Treatment	Dose (kg ha-1)	Time of application
1	Oxyfluorfen <i>fb</i> one hand weeding	0.10	Post emergence and hand weeding at 60 (DAT)
2	Oxyfluorfen	0.20	Post - emergence (40 DAT)
3.	Oxyfluorfen	0.30	Post - emergence (40 DAT)
4.	Pendimethalin	1.5	Pre emergence
5.	Quizalofop <i>fb</i> one hand weeding	0.06	Post- emergence (40 DAT) and hand weeding at 60 (DAT)
6.	Two hand weeding	–	(40 and 60 DAT)
7.	Farmer practice (three hand weeding)	–	30, 60 and 90 (DAT)
8.	Weed free	–	20 DAT onwards
9.	Weedy check	–	–

3. Results and Discussion

3.1 Weed dry weight

At harvest of the crop oxyfluorfen @ 0.30 kg ha⁻¹ recorded significantly less dry weight of weeds (1.32 g m⁻²). The minimum dry weight of weeds was observed in weed free (1.00 g m⁻²) treatment. The maximum dry weight of weed was recorded in weedy check (15.87 g m⁻²) treatment. The minimum dry weight of weeds at all the stages of plant growth was recorded when oxyfluorfen @ 0.30 kg ha⁻¹ was applied among herbicidal treatments and in weed free treatment among hand weeding. Rahman *et al.* (2012) [9] also found manual weeding very effective in crop to control weed biomass. The results are also in agreement with the finding of Nandal and Ravinder (2002) [6]. According to Patel *et al.* (2012) [7] dry weight of weeds may be due to the increased weed population and continuous growth and may also be due to the higher amount of nutrient uptake.

3.2 Plant Height

All the weed control treatments recorded maximum plant height when compared with weedy check. The maximum plant height (55.88 cm) was recorded in weed free treatment followed by two hand weeding (37.95 cm). Among herbicidal treatments oxyfluorfen @ 0.30 kg ha⁻¹ recorded the maximum plant height (53.35 cm) and it was significantly higher from

all other herbicidal treatments. The minimum plant height (13.69 cm) was recorded in weedy check treatment. The same results were found by Anisuzzaman *et al.* (2009) [2]. According to Patel *et al.* (2012) [7] it was due to favorable environment in the root zone resulting in absorption of more water and nutrient from soil and good control of weed competition throughout the different growth stages of the crop.

3.3 Plant population

Plant population recorded at harvest varied significantly with different weed control treatments. The numbers of plants were significantly higher in all the treatments when compared with weedy check. The maximum plant stand was recorded in weed free plot (276.33 per plot) and significantly higher among manual as well as herbicidal treatments. Oxyfluorfen @ 0.30 kg ha⁻¹ (258.33 per plot) recorded maximum plant stand which was statistically at par with oxyfluorfen @ 0.20 kg ha⁻¹ (251.67 per plot). The minimum plant stand was recorded in weedy check (204.33 per plot). This is may be due to the presence of higher number of weeds which competed with onion bulb for light, water, space and nutrients. Weed free treatment was responsible for better plant growth and development which resulted in better plant stand. Similar results were obtained by Jilani *et al.* (2007) [4].

Table 2: Effect of treatments on weed dry weight, plant height, plant population and bulb yield

Treatments	weed dry weight (g m ⁻²)	Plant Height (cm)	Plant population	Bulb yield q ha ⁻¹
Oxyfluorfen fb one HW (0.10 kg ha ⁻¹)	3.42	26.33	232.67	148.61
Oxyfluorfen (0.20 kg ha ⁻¹)	1.90	41.88	251.67	226.61
Oxyfluorfen (0.30 kg ha ⁻¹)	1.32	53.35	258.33	258.33
Pendimethalin (1.5 kg ha ⁻¹)	3.78	17.31	228.33	174.44
Quizalofop fb one HW (0.06 kg ha ⁻¹)	4.52	23.58	231.67	128.33
Two hand weeding	2.85	37.95	230.33	120.94
Farmer practice (Three HW)	1.35	33.61	241.33	218.89
Weed free	1.00	55.88	276.33	272.22
Weedy check	15.87	13.69	204.33	84.17
CD (P=0.05)	0.54	3.47	11.21	17.94

3.4 Bulb Yield (q ha⁻¹)

The yield data (q ha⁻¹) was recorded during 2014. The yield was significantly higher in all the treatments as compared to weedy check. In weed free treatment maximum bulb yield (272.22 q ha⁻¹) was recorded which was significantly higher from other treatments. Oxyfluorfen @ 0.30 kg ha⁻¹ significantly recorded maximum yield (258.33 q ha⁻¹) and significantly higher from other treatments. The minimum bulb yield was recorded in weedy check treatment (84.17 q ha⁻¹). This might be due conservation of nutrients, increased soil temperature, prevention of water evaporation from soil and sustained adequate soil moisture at outer soil surface, where most onion root occur 0 to 30 cm. The refore, searching for moisture to further soil depths is not required which in turn source sink assimilate for the vegetative growth and yield.

4. References

1. Anonymous, 2014 .www.nhb.gov.in
2. Anisuzzaman M, Ashrafuzzaman M, Mohd Razi Ismail, Uddin MK, Rahim MA. Planting time and mulching effect on onion development and seed production. African Journal of Biotechnology. 2009; 8(3):412-416
3. Kashyap S. Studies on integrated weed management in capsicum (*Capsicum annuum* L.). 54p. M.Sc. Thesis. Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan, H.P. India, 2010.
4. Jilani MS, Ramzan M, Waseem K. Impact of weed management practices on growth and yield of some local genotypes of onion. Pakistan Journal of Weed Science Research. 2007; 13(3/4):191-198
5. Khatam A, Ahmad W, Khan MZ. Perception of farmers regarding effect of various weed management practices in onion crop. Pakistan Journal of Weed Science Research. 2012; 18(4):553-559
6. Nandal TR, Ravinder Singh. Integrated weed management in onion (*Allium cepa* L.) under Himachal Pradesh condition. Indian Journal of Weed Science. 2002; 34(1/2):72-75
7. Patel TU, Patel CL, Patel DD, Thanki JD, Arvadia MK, Vaidya HB. Performance of onion under weed and fertilizer management. Indian Journal of Weed Science 2012; 44(3):151-158
8. Pramanick M, Roy A, Debnath S, Dwary A. An analysis on growth, productivity, and economics of onion under different coloured polythene mulching. Horticulture Journal. 2007; 20(1):17-18
9. Rahman H, Ullah K, Sadiq M, Javaria S, Ullah I, Khan MA. Relationship between manual weeds removal timings and onion yield. Pakistan Journal of Weed Science Research. 2012; 18(2):201-207
10. Ramalingam SP, Chinnagounder C, Perumal M,

Palanisamy MA. Evaluation of New Formulation of Oxyfluorfen (23.5% EC) for Weed Control Efficacy and Bulb Yield in Onion. American Journal of Plant Sciences 2013; 4:890-895

11. Singh SJ, Sinha KK, Mishra SS, Thakur SS, Choudhary NK. Studies on weed management in onion. Indian Journal of Weed Science 1992; 24:6-10.
12. Uygur S, Gurbuz R, Uygur FN. Weeds of onion fields and effects of some herbicides on weeds in Cukurova region, Turkey. African Journal of Biotechnology. 2010; 9(42):7037-7042.
13. Vavilov NI. Origin, variation, immunity and breeding of cultivated plants. Chron Bot. 1951; 13(1-6):1-364.