



E-ISSN: 2278-4136  
P-ISSN: 2349-8234  
JPP 2019; 8(3): 01-05  
Received: 01-03-2019  
Accepted: 04-04-2019

**Subhash Mishra**

Department of Genetics and  
Plant Breeding NDU&T  
Kumarganj, Ayodhya,  
Uttar Pradesh India

**OP Verma**

Department of Genetics and  
Plant Breeding NDU&T  
Kumarganj, Ayodhya,  
Uttar Pradesh India

**Ashish**

Department of Genetics and  
Plant Breeding NDU&T  
Kumarganj, Ayodhya,  
Uttar Pradesh India

**SP Srivastava**

Department of Genetics and  
Plant Breeding NDU&T  
Kumarganj, Ayodhya,  
Uttar Pradesh India

**KP Singh**

Department of Genetics and  
Plant Breeding NDU&T  
Kumarganj, Ayodhya,  
Uttar Pradesh India

**Correspondence****Subhash Mishra**

Department of Genetics and  
Plant Breeding NDU&T  
Kumarganj, Ayodhya,  
Uttar Pradesh India

## Identifying superior parents and hybrids for certain quantitative traits in rice (*Oryza sativa* L.) under sodic soils

Subhash Mishra, OP Verma, Ashish, SP Srivastava and KP Singh

**Abstract**

The present investigation entitled “Studies on genetic variability, combining ability and heterosis in aromatic and non-aromatic rice (*Oryza sativa* L.) unersodic soil. The present study was based on two related experiments, namely, Germplasm evaluation (Experiment-I) and Combining ability (Experiment-II). These experiments were conducted at Main experimental farm, of the N.D. University of Agriculture & Technology, Narendra Nagar (Kumarganj), Ayodhya, during *Kharif*, 2015 and 2016. The combining ability experiment was based on evaluation of a line x tester set of 69 hybrids ( $F_1$ 's) and their 26 parents along with two checks for seven characters under sodic soil condition in randomized complete block design with three replications during *Kharif*, 2016. The 69  $F_1$ 's were generated by crossing 23 lines with 3 testers during *Kharif*, 2015.

Analysis of variance for combining ability revealed non-significant mean squares due to testers and lines for all the characters under study except significant mean squares observed for testers for plant height. This indicated less importance of general combining ability and additive gene effects in expression of out of 7 characters. The mean squares due to lines x testers interactions, indicating importance of specific combining ability and non-additive gene effects, were found to be highly significant for all the seven characters under study. Twenty-three out of sixty-nine crosses emerged with positive and significant sca effects for grain yield/plant. The five best crosses were Pusa Bas.-1 x Jaya, NDRK-50046 x Jaya, NDRK-50055 x Jaya, NDRK-50054 x CSR-10 and NDRK-50063 x CSR-10 which showed significant and positive sca effects for grain yield/plant as well as some other yield components.

**Keywords:** Rice, combining ability, general and specific combining ability

**Introduction**

Rice (*Oryza sativa* L.  $2n = 24$ ), belongs to the family gramineae (Poaceae). Rice is a staple food of developing world as well as majority of Asian countries. More than 90 per cent of the world's rice is grown and consumed in Asia (rice bowl of the world), where 16 per cent of the earth's people and two third of world's poor live (Khush and Virk 2000) [4]. Globally, it stands first in rice area and second in rice production, after China. In India, rice is cultivated over an area of about 42.77 million hectares with an annual production of about 106.54 million tonnes with an average productivity of 2490 kg per hectare (Anonymous, 2014) [2]. Uttar Pradesh is an important rice growing state in the country. The area and production of rice in this state is about 5.95 million hectares and 13.53 million tonnes, respectively, with productivity of 2273 kg per hectare (Anonymous, 2014) [1].

An understanding of the nature and magnitude of combining ability would, therefore, be of great importance in breeding high yielding rice genotypes. The line x tester analysis has been found to be helpful to plant breeders in identifying most promising parents and hybrids for further exploitation in breeding programmes.

**Materials and Methods**

The technical programme of the present investigation will be based on two experiments as follows:

**Genetic variability (Experiment-I):** This experiment will involve evaluation of 115 elite aromatic and non-aromatic rice lines along with 3 checks *viz.* Sarjoo 52, FL 478 and CSR 43. The 115 rice lines along with three checks will be evaluated in augmented design during *Kharif*, 2016. The experimental plot will be sub-divided in to 10 blocks of 13 plots each. The three checks will be allotted randomly to three plots in each block, while remaining 10 plots in a block will be used for accommodating the unreplicated test genotypes.

**Combining ability analysis (Experiment-II):** A line x tester set of 69 hybrids ( $F_1$ 's) will be derived by crossing 23 rice genotypes /varieties as lines (females) with three testers (males) viz., Jaya, BPT 5204 and CSR 10. The twenty three lines are NDRK 50063, Narendra Usar 2009, NDRK 50051, Narendra 6093, Narendra 6330, Pusa Basmati 1, Improved Pusa Basmati -1, Narendra 359, NDRK 50054, NDRK 50060, NDRK50046, NDRK50058, NDRK 50055, NDRK 50052, IRSSTN 150, IR-14 T-102, IR-14 T-105, N 80, Usar 1, Narendra 2026, N-3112-1, Narendra Shushk Samrat and Narendra Parag. The 69  $F_1$ s along with their parents and check varieties (Narendra Usar Dhan 3 and IR 28), will be evaluated in randomized complete block design with three replications during *Kharif*2016. Single row plots of 3m length will be used in both the experiments for transplanting single seedling per hill for each genotype (treatment) following intra-row and inter-row spacing of 15 cm and 20 cm, respectively. Recommended cultural practices will be followed to raise a good crop in both the experiments.

## Results

The character wise results of specific combining ability effect of sixty-nine crosses are presented in Table: 2 and described here under;

The understanding of inheritance of various characters and identification of superior parents are important pre-requisites for launching an effective and efficient breeding programme (Dhillon, 1975). Combining ability analysis is useful technique for understanding of genetic worth of parents and their crosses for further exploitation in breeding programmes. Among the various techniques of combining ability analysis, line x tester analysis (Kempthorne, 1957) has been widely utilized for screening of germplasm to identify valuable donor parents and promising crosses in many crops including rice (Satheesh *et al.*, 2010; Saidaiah *et al.*, 2011; Devi *et al.*, 2017) [6, 3].

The significant and positive GCA effects for grain yield/plant were exhibited by ten lines, NDRK-50052 (1.79), Pusa Basmati-1, NDRK-50060 (1.24), NDRK-50055 (1.23), Usar-1(1.07), NDRK-50046 (1.06), NDRK-50058 (0.94), NDRK-50063 (0.51), IR-92831-22-BAY-2-1 (0.36) and NDR-359 (0.29) The parent with highest GCA effects for grain yield, NDRK 50052, also showed significant GCA effects in desirable direction for days to 50% flowering, flag leaf area and chlorophyll content. Tester CSR-10 showed good general combiner for grain yield/plant and exhibited significant and positive GCA effect for plant height and chlorophyll content under sodic soil.

The eleven parents showing positive and significant GCA effects for grain yield and other important traits as mentioned above may serve as valuable parents for hybridization

programme or multiple crossing programme for obtaining high yielding hybrid varieties or transgressive segregants for developing pure line varieties of rice sodic soil. It is evident from Table 2, that most of the lines showing significant positive GCA effects for grain yield per plant also exhibited positive and significant GCA effects for some of the important yield components traits. This indicated that the significant GCA effects for grain yield in positive direction resulted from similar GCA effects of some other yield components suggesting that the combining ability for grain yield was influenced by the combining ability of its components. Therefore, simultaneous improvement in important yield components and other associated traits along with grain yield may be better approach for raising yield potential in rice under sodic soil. These findings are similar to those of Verma and Srivastava (2004) [8].

## Specific combining ability effects

The specific combining ability (SCA) effects, which are supposed to be manifestation of non-additive components of genetic variance, are highly valuable for discrimination of crosses for their genetic worth as breeding materials. The estimates of SCA effects of sixty-nine crosses for thirteen characters are given in Table 3. Seventeen out of sixty-nine crosses emerged with positive and significant SCA effects for grain yield/plant. The top ten crosses were Pusa Bas.-1 x Jaya, NDRK-50046 x Jaya, NDRK-50055 x Jaya, NDRK-50054 x CSR-10, NDRK-50063 x CSR-10, IR 14 T 102 x BPT 5204, NDRK 50058 x CSR 10, Usar 1 x CSR 10, IR 14 T 105 x BPT 5204 and NDRK 50060 x CSR 10 showed significant and positive SCA effects for grain yield/plant as well as some other yield components. The cross having highest positive and significant SCA effects for grain yield/plant Pusa Basmati-1 x Jaya also recorded significant SCA effects in desirable direction for plant height, flag leaf area and chlorophyll content. The second ranking cross for higher positive and significant SCA effect for grain yield/plant, NDRK-50046 x Jaya showed significant and desirable SCA effects for panicle bearing tillers/plant and panicle length,. Similarly, remaining fifteen crosses having significant and positive SCA effects for grain yield/plant also possessed significant SCA effects in desirable direction for some other characters also.

The foregoing observation clearly indicated that there was no particular relationship between positive and significant SCA effects of crosses with GCA effects of their parents for the characters under study. Verma and Srivastava, (2004) +, Singh and Kumar (2004) [7] and Devi *et al.*, (2017) [3] have also found that crosses having high order positive SCA effects for grain yield resulted from parents having high x high as well as high x low GCA effects for grain yield.

**Table 1:** Analysis of variance for combining ability following line  $\times$  tester mating design for 13 characters in rice under sodic soil

Characters d.f.	Sources of variation				
	Replications	Lines	Testers	Lines $\times$ Testers	Error
	2	22	2	44	136
Days to 50% flowering	0.83	7.0	7.36	14.33**	0.38
Plant height (cm)	0.76	13.94	20.89	30.40**	0.29
Flag leaf area (cm <sup>2</sup> )	0.04	5.19	35.94**	4.63**	0.17
Chlorophyll content (SPAD Value)	0.00	5.29	2.33	4.64**	0.05
Panicle bearing tillers / plant	0.01	1.31	8.90*	1.78**	0.21
Panicle length (cm)	0.14	15.38	1.28	10.07**	0.40
Grain yield / plant (g)	0.04	8.56	23.70*	5.37**	0.05

\*, \*\* Significant at 5% and 1% probability levels, respectively

**Table 2:** Estimates of general combining ability (GCA) effects of parents (lines and testers) for 13 characters in rice under sodic soil

S. No.	Line	Days to 50% flowering	Plant height (cm)	Flag leaf area (m <sup>2</sup> )	Chlorophyll Content (SPAD Value)	Panicle bearing tillers / plant	Panicle length (cm)	Grain yield / plant (g)
1	NDRK - 50055	1.04*	-0.30	-2.48**	-0.59**	-0.98**	-0.94**	1.23**
2	Imp. Pusa Bas. -1	1.37**	0.08	-0.71**	0.85**	-0.44**	-1.64**	-0.33**
3	Narendra - 359	-0.30	-1.93**	-1.44**	0.23**	0.45**	2.36**	0.29**
4	NDRK - 50051	-0.18	0.92**	0.07	0.09	0.55**	-2.62**	-0.85**
5	Narendra - 6330	-1.30**	-0.95**	0.06	-0.28**	0.32*	0.07	-1.49**
6	NDRK - 50063	0.48	-1.47**	0.15	0.00	0.65**	0.13	0.51**
7	IR - 14 T -102	-0.63	0.30	0.39**	-0.70**	0.40**	0.66**	-1.22**
8	N. Usar - 2009	2.04**	-0.14	0.41**	0.01	0.14	0.45*	-0.76**
9	IR - 14 T -105	0.82*	-0.66**	0.49**	-0.91**	-0.03	2.20**	-0.54**
10	NDRK - 50054	-0.96*	-1.20**	0.82**	0.94**	-0.01	1.87**	-0.25**
11	NDR - 2026	-0.63	1.78**	-0.27*	-0.05	0.06	1.82**	-1.53**
12	Narendra - 3112 - 1	-1.52**	-2.55**	0.52**	-0.82**	0.18	-0.64**	-0.39**
13	NDRK - 50046	-0.63	0.14	0.19	0.46**	-0.38**	1.13**	1.06**
14	Usar - 1	-0.63	1.87**	0.81**	-0.36**	-0.13	0.62**	1.07**
15	N - 80	0.48	1.07**	0.23	0.02	-0.49**	-1.29**	-0.74**
16	IR-92831-22-BAY 2-1	-0.30	-2.17**	0.33**	-1.19**	-0.10	-0.90**	0.36**
17	N. Parag	0.59	0.95**	0.58**	0.63**	0.06	0.08	-0.79**
18	Pusa Bas. - 1	0.71	0.78**	-0.32*	0.16*	0.36*	-0.29	1.24**
19	NDRK - 50058	0.26	0.40	-0.88**	1.50**	0.06	-0.71**	0.94**
20	N. ShushkSamrat	0.48	0.01	0.28*	-1.27**	-0.38**	-0.94**	-0.45**
21	NDRK - 50052	-0.85*	0.60**	0.25*	1.52**	0.09	-0.57**	1.79**
22	N - 6093	0.15	1.54**	0.32*	-0.54**	-0.35*	-1.55**	-0.39**
23	NDRK - 50060	-0.52	0.93**	0.22	-0.44**	-0.02	0.71**	1.24**
SE (gi) Lines		0.79	0.41	0.24	0.16	0.27	0.42	0.15
SE(gi - gj)		1.12	0.58	0.35	0.22	0.39	0.60	0.22
Tester								
1	CSR - 10	0.20	-0.63**	0.08	0.16**	0.07	0.08	0.68**
2	BPT - 5204	-0.38**	0.30**	0.68**	-0.20**	0.32**	-0.16*	-0.37**
3	Jaya	0.17	0.34**	-0.76**	0.04	-0.39**	0.08	-0.31**
SE(gi) Tester		0.29	0.15	0.09	0.06	0.10	0.15	0.06
SE(gi - gj)		0.40	0.21	0.12	0.08	0.14	0.22	0.08

**Table 3:** Estimates of specific combining ability (SCA) effects of crosses for 7 characters in rice under sodic soil

S. No.	Crosses	Days to 50% flowering	Plant height (cm)	Flag leaf area (cm <sup>2</sup> )	Chlorophyll content (SPAD Value)	Panicle bearing tillers / plant	Panicle length (cm)	Grain yield / plant (g)
1	NDRK-50055 X CSR-10	0.69	0.28	-1.91**	-1.21**	-0.65**	0.30	-1.38**
2	NDRK-50055 X BPT-5204	0.27	1.32**	-2.03**	0.15	-0.43	-1.53**	-0.87**
3	NDRK-50055 X Jaya	-0.95	-1.59**	3.94**	1.05**	1.08**	1.23**	2.24**
4	Imp. Pusa Bas. 1 X CSR-10	0.69	1.03**	1.42**	-0.19	0.92**	1.63**	-0.92**
5	Imp. Pusa Bas. 1 X BPT-5204	-0.73	4.00**	-0.48*	-0.02	0.00	-0.17	0.76**
6	Imp. Pusa Bas1 X Jaya	0.05	-5.04**	-0.94**	0.21	-0.92**	-1.47**	0.16
7	Narendra-359 X CSR-10	0.02	-1.15**	1.25**	1.17**	1.27**	-0.23	1.26**
8	Narendra-359 X BPT-5204	1.60*	-0.75*	-2.11**	-1.37**	-0.49*	1.27**	-0.60**
9	Narendra-359 X Jaya	-1.62*	1.91**	0.86**	0.20	-0.78**	-1.04**	-0.66**
10	NDRK-50051 X CSR-10	1.24	3.42**	-0.33	-0.75**	0.90**	0.01	-0.46**
11	NDRK-50051 X BPT-5204	-1.51*	-1.87**	-1.12**	-1.36**	0.65**	-0.55	1.11**
12	NDRK-50051 X Jaya	0.27	-1.55**	1.45**	2.11**	-1.54**	0.54	-0.65**
13	Narendra-6330 X CSR-10	2.02**	-4.00**	1.45**	-0.52**	-0.21	-0.54	0.11
14	Narendra-6330 X BPT-5204	-1.07	0.17	-1.04**	1.58**	-0.26	1.06**	0.12
15	Narendra-6330 X Jaya	-0.95	3.83**	-0.41	-1.06**	0.48*	-0.51	-0.24
16	NDRK-50063 X CSR-10	-1.43*	-2.31**	1.46**	-0.43**	0.97**	-0.14	1.98**
17	NDRK-50063 X BPT-5204	1.82**	1.79**	-0.97**	0.83**	-0.19	0.92*	-0.14
18	NDRK-50063 X Jaya	-0.40	0.52	-0.50*	-0.40**	-0.78**	-0.78*	-1.84**
19	IR-14 T-102 X CSR-10	-1.31	2.68**	1.22**	2.17**	-0.16	1.37**	-1.56**
20	IR-14 T-102 X BPT-5204	-2.07**	0.78*	-0.48*	-0.47**	-0.61*	-0.73	1.81**
21	IR-14 T-102 X Jaya	3.38**	-3.46**	-0.74**	-1.70**	0.77**	-0.64	-0.25
22	N. Usar-2009 X CSR-10	-0.31	1.22**	1.11**	-0.54**	0.11	-1.35**	0.28*
23	N. Usar-2009 X BPT-5204	-0.40	1.19**	-0.05	2.45**	0.16	1.81**	0.29*
24	N. Usar-2009 X Jaya	0.71	-2.42**	-1.05**	-1.91**	-0.27	-0.46	-0.57**
25	IR-14 T-105 X CSR-10	2.24**	3.01**	0.88**	0.91**	0.28	1.79**	-0.61**
26	IR-14 T-105 X BPT-5204	-3.51**	-0.98**	-0.04	-1.02**	-0.18	0.12	1.33**
27	IR-14 T-105 X Jaya	1.27	-2.03**	-0.84**	0.11	-0.10	-1.91**	-0.73**
28	NDRK-50054 X CSR-10	-0.31	-3.65**	-0.57**	1.06**	0.35	2.06**	2.04**
29	NDRK-50054 X BPT-5204	-1.40*	3.32**	0.03	-0.38**	-0.23	-1.48**	-1.52**

30	NDRK-50054 X Jaya	1.71*	0.34	0.54*	-0.68**	-0.12	-0.58	-0.51**
31	NDR-2026 X CSR-10	-2.65**	1.30**	-0.88**	1.11**	0.49*	2.37**	0.35*
32	NDR-2026 X BPT-5204	0.27	-3.63**	1.22**	-2.09**	-0.30	-1.47**	0.09
33	NDR-2026 X Jaya	2.38**	2.33**	-0.34	0.98**	-0.19	-0.90*	-0.44**
34	Narendra-3112-1 X CSR-10	-0.76	5.63**	-0.34	-1.52**	0.73**	-1.70**	-0.89**
35	Narendra-3112-1 X BPT-5204	-0.85	-1.30**	0.40	0.01	-0.42	1.83**	0.16
36	Narendra-3112-1 X Jaya	1.60*	-4.34**	-0.06	1.51**	-0.31	-0.14	0.73**
37	NDRK-50046 X CSR-10	2.69**	4.95**	0.02	0.69**	-0.71**	-1.88**	-1.68**
38	NDRK-50046 X BPT-5204	-0.73	-0.98**	0.69**	-0.01	0.14	-1.18**	-0.63**
39	NDRK-50046 X Jaya	-1.95**	-3.96**	-0.71**	-0.68**	0.58*	3.05**	2.31**
40	Usar-1 X CSR-10	0.35	2.91**	-0.49*	1.79**	-0.96**	-4.43**	1.35**
41	Usar-1 X BPT-5204	-1.73*	-3.72**	0.21	-0.68**	0.32	3.83**	-0.78**
42	Usar-1 X Jaya	1.38*	0.81*	0.28	-1.11**	0.63**	0.60	-0.57**
43	N-80 X CSR-10	-2.43**	0.01	0.38	0.18	-0.60*	-1.49**	0.89**
44	N-80 X BPT-5204	2.15**	2.95**	0.32	0.74**	0.25	1.11**	-1.17**
45	N-80 X Jaya	0.27	-2.96**	-0.71**	-0.92**	0.36	0.37	0.27*
46	IR-92831-22-BAY 2-1 X CSR-10	3.35**	-2.61**	-0.62**	-0.01	-0.99**	-0.14	-0.08
47	IR-92831-22-BAY 2-1 X BPT-5204	-0.40	3.33**	0.52*	-0.15	-0.14	-0.94*	0.10
48	IR-92831-22-BAY 2-1 X Jaya	-2.95**	-0.72*	0.09	0.15	1.13**	1.09**	-0.03
49	N. Parag X CSR-10	0.46	-0.87*	0.06	-1.13**	0.49*	3.78**	0.28*
50	N. Parag X BPT-5204	-1.96**	-1.80**	0.17	1.67**	-0.30	-2.02**	0.56**
51	N. Parag X Jaya	1.49*	2.66**	-0.23	-0.54**	-0.19	-1.76**	-0.84**
52	Pusa Bas.-1 X CSR-10	0.35	0.30	-1.11**	-0.60**	-1.45**	0.08	-1.72**
53	Pusa Bas.-1 X BPT-5204	3.60**	-2.63**	0.67**	-0.07	1.20**	-0.55	-1.04**
54	Pusa Bas.-1 X Jaya	-3.95**	2.33**	0.44*	0.66**	0.24	0.47	2.76**
55	NDRK-50058 X CSR-10	-1.20	-3.32**	-1.14**	-0.36**	0.49*	-1.60**	1.75**
56	NDRK-50058 X BPT-5204	1.38*	0.75*	0.87**	-0.27	-0.30	1.13**	-1.44**
57	NDRK-50058 X Jaya	-0.17	2.57**	0.27	0.63**	-0.19	0.46	-0.30*
58	N. ShushkSamrat X CSR-10	1.57*	-2.00**	0.39	-0.70**	-0.38	-0.07	-0.83**
59	N. ShushkSamrat X BPT-5204	-1.85**	-0.86*	0.53*	0.37**	0.14	-1.00**	0.61**
60	N. ShushkSamrat X Jaya	0.27	2.86**	-0.93**	0.33*	0.24	1.06**	0.22
61	NDRK-50052 X CSR-10	-0.43	-1.49**	0.19	-0.45**	-0.88**	-0.44	-1.34**
62	NDRK-50052 X BPT-5204	2.49**	-1.68**	0.57**	0.08	1.10**	0.39	1.83**
63	NDRK-50052 X Jaya	-2.06**	3.17**	-0.76**	0.38**	-0.22	0.05	-0.49**
64	N-6093 X CSR-10	-3.43**	-2.25**	-0.71**	-0.86**	-0.33	-1.49**	-0.12
65	N-6093 X BPT-5204	2.49**	-0.32	0.57**	-0.50**	0.11	-0.25	0.02
66	N-6093 X Jaya	0.94	2.57**	0.14	1.36**	0.22	1.74**	0.10
67	NDRK-50060 X CSR-10	-1.43*	-3.11**	-1.74**	0.17	0.33	2.11**	1.31**
68	NDRK-50060 X BPT-5204	2.15**	0.93*	1.53**	0.50**	-0.22	-1.62**	-0.61**
69	NDRK-50060 X Jaya	-0.73	2.18**	0.20	-0.67**	-0.11	-0.49	-0.70**
	SE (S <sub>ij</sub> )	1.37	0.70	0.42	0.27	0.48	0.74	0.27
	SE (S <sub>ij</sub> - S <sub>kl</sub> )	1.93	1.00	0.60	0.38	0.67	1.05	0.38

\*, \*\* significant at 5 and 1 per cent probability levels, respectively

## Discussion

The analysis of variance for combining ability revealed non-significant mean squares due to testers and lines for all the characters under study except significant mean squares observed for testers for plant height. This indicated less importance of general combining ability and additive gene effects in expression of out of 13 characters. The mean squares due to lines x testers interactions, indicating importance of specific combining ability and non-additive gene effects, were found to be highly significant for all the thirteen characters under study.

The estimates of SCA variance were higher than the corresponding estimates of gca variance for all the traits. On the basis of general combining ability effects, NDRK 50052 and Pusa Basmati 1, NDRK 50060 showed desirable and significant gca effects for grain yield/plant and some of the yield contributing traits to emerge as valuable donor parents for hybridization programme for obtaining high yielding hybrid varieties or transgressive segregants for developing pure line varieties of rice under sodic soil.

Twenty-three out of sixty-nine crosses emerged with positive and significant SCA effects for grain yield/plant. The top ten crosses were Pusa Bas.-1 x Jaya, NDRK-50046 x Jaya,

NDRK-50055 x Jaya, NDRK-50054 x CSR-10, NDRK-50063 x CSR-10, IR 14 T 102 x BPT 5204, NDRK 50058 x CSR 10, Usar 1 x CSR 10, IT 14 T105 x BPT 5204 and NDRK 50060 x CSR 10 which showed significant and positive SCA effects for grain yield/plant as well as some other yield components. The twenty-three crosses having positive and desirable SCA effects for grain yield and some of its component traits merit attention for exploitation as hybrid varieties in breeding materials for developing superior aerobic rice genotypes under salt affected soil.

The twenty-three crosses having significant and positive SCA effects for grain yield per plant also showed desirable and significant SCA effects for some other characters, most commonly biological yield/plant, 1000-grain weight, harvest-index, flag leaf area and spikelet fertility suggested that manifestation of SCA effects for grain yield is related with higher SCA effects for important yield components. The crosses exhibiting high order significant and desirable SCA effects for different characters involved parents having all types of combinations of gca effects such as high x high (H x H), high x average (H x A), high x low (H x L), average x average (A x A), average x low (A x L) and low x low (L x L) general combiner parents indicating thereby lack of any

relationship between positive and significant SCA effects of crosses with gca effects of their parents for the characters under study.

### References

1. Anonymous. FAO statistical data base on agriculture, [http// apps.FAO.org](http://apps.FAO.org), 2014.
2. Anonymous. U.P. Directorate of Agricultural Statistics Reports, 2014.
3. Devi Archana, Kumari Preeti, Dwivedi Ranjan, Dwivedi Saket, Verma OP, Singh PK. *et al.* Gene action and combining ability analysis for yield and yield contributing traits in rice (*Oryza sativa* L.) over environment, J of Pharmacognosy and Phytochemistry. 2017; 6(3):662-671.
4. Khush GS, Virk PS. Rice breeding achievements and future strategies. Crop Improvement. 2000; 27(2):115-144.
5. Saidaiah P, Sudheer Kumar S, Ramesha MS, Combining ability studies for development of new hybrids in rice over environments. J of Agric. Sci. 2010; 2(2):225-233.
6. Satheesh KP, Saravanan K, Sabesan T. Combining ability for yield & yield contributing characters in rice (*Oryza sativa* L.). Ele. J of Plant Breeding. 2010; 1(5):1290-1293.
7. Singh NK, Kumar Anand. Combining ability analysis to identify suitable parents for heterotic rice hybrid breeding. Int. Rice Res. Notes. 2004; 29(1):21-22.
8. Verma OP, Srivastava HK. Genetic component and combining ability analysis in relation to heterosis for yiel and its associated traits using three divarce rice growing ecosystem. Field Crop Rse. 2004; 88:91-102.