



E-ISSN: 2278-4136

P-ISSN: 2349-8234

JPP 2019; 8(5): 2524-2527

Received: 22-07-2019

Accepted: 28-08-2019

Dr. Vishal Bharat Babar

PG Research Department of
Pharmaceutical Chemistry,
Dattakala College of Pharmacy,
Swami-Chincholi, Tal: Daund,
Dist: Pune, Maharashtra, India

Prajwala Rajkumar Khapale

Department of Pharmacology
and Toxicology, Dattakala
College of Pharmacy, Swami-
Chincholi, Tal: Daund, Dist:
Pune, Maharashtra, India

Sudarshan Narayan Nagarale

PG Research Department of
Pharmaceutical Chemistry,
Dattakala College of Pharmacy,
Swami-Chincholi, Tal: Daund,
Dist: Pune, Maharashtra, India

Corresponding Author:**Dr. Vishal Bharat Babar**

PG Research Department of
Pharmaceutical Chemistry,
Dattakala College of Pharmacy,
Swami-Chincholi, Tal: Daund,
Dist: Pune, Maharashtra, India

Preliminary investigation and in-vitro anticancer activity of *Lantana camara* L. (Verbenaceae)

Dr. Vishal Bharat Babar, Prajwala Rajkumar Khapale and Sudarshan Narayan Nagarale

Abstract

The present study is the evaluation of anticancer activity of *Lantana camara* L. stem extracts and also an investigation of the phytoconstituents from the different extracts of the plant. Chloroform, ethyl acetate, methanol and aqueous extracts of *Lantana camara* L. stem were tested for their anticancer activity on the cancer cell lines HL-60 (Leukemia) at various concentrations. The anticancer effect of the extracts on cell inhibition was studied using Sulphorhodamine B assay. Only chloroform, ethyl acetate, and methanolic extracts showed the dose-dependent anticancer effect on leukemia cancer cells. Phytochemical studies of the plant extracts showed the presence of flavonoid, alkaloids, saponins, tannins, and triterpenes. The result revealed that chloroform, ethyl acetate and methanolic extracts of *Lantana camara* L. contains important chemical constituents, which can be further used in the treatment of leukemia.

Keywords: *Lantana camara* L., Anticancer, Lukemia, Sulphorhodamine B assay, HL-60 etc.

Introduction

Cancer is one of the most life-threatening diseases and the leading causes of death all over the world. According to cancer statistics in 2013, stomach and liver cancer are the most common in Asia, and both are associated with high mortality rates, while bladder cancer is the most common in the USA. Colorectal and breast cancers have high incidence rates in all countries [1]. According to the estimates of the WHO, more than 80% of people in developing countries depend on traditional medicine for their primary health needs [2]. The plant kingdom has provided a variety of medicines for cancer treatment, currently, over 60% of the drugs are derived in one or other way from natural source including plant, marine organism and micro-organism [3]. Over the past decade, herbal medicines have been accepted universally, and they have an impact on both world health and international trade. Hence, medicinal plants continue to play an important role in the healthcare system of a large number of the world's population. Traditional medicine is widely used in India. The National Cancer Institute collected about 35,000 plant samples from 20 different countries and has screened around 114,000 extracts for anticancer activity [4].

Treatment by herbal medicines may have some advantages over treatment by single purified chemicals; as herbal medicine are the mixtures of more therapeutic or preventive components, and so might have more activity than single products alone [5]. The antioxidant and anti-tumor effects of extracts from various herbs and medicinal plants have been proved experimentally and clinically. Several in-vitro or in-vivo studies have proved the anticancer potential of the extracts from several medicinal plants [6]. A diverse number of anticancer chemical constituents have been isolated from natural sources viz. Camptothecin, vincristine, vinblastine, taxol, podophyllotoxin, etc. Further modification of these compounds leads to the development of potent anticancer agents like Topotecan, Irinotecan, and etoposide teniposide, etc. [7]

Lantana camara L. is commonly known as pigeon berry belongs to the family Verbenaceae. It is shrubs or herbs usually 1 to 3 m in height [8]. The plant is not browsed by cattle and is believed to be poisonous [9]. Ethyl acetate and aqueous extracts of leaves showed significant antimalarial activity when administered to mice 10. The fruits are used in the treatment of malaria and intestinal worms [11]. The leaves are used in the treatment of abscess [12]. From the genus *Lantana camara* several iridoid glycosides as camarantosides I, II, III, IV, and lamiide was isolated [13-14]. Flavonoids and C-alkylated flavonoids [15-16] and some alkaloids [17] were isolated. Based on the literature survey, it is evident that no work has been carried out on the evaluation of the anticancer property of stem extracts on the selected cell lines. Hence in this present study, the anticancer activity of *Lantana camara* L. stem extracts was assessed by investigating the inhibition of cell growth of HL-60 (Leukemia) at various concentrations.

Materials and Methods

Plant Materials and Reagents

The stems of *Lantana camara* L. were picked from the rural area of Bhigwan, Pune District, Maharashtra. The specimen sample was taxonomically identified and authenticated by the Department of Botany, Dattakala School and Jr. College, Swami-Chincholi, where a voucher specimen was deposited (authentication no. DKJrCBL No. 2019119) this medicinal material has a scientific name of *Lantana camara* L. All the reagents and chemicals were purchased from Merck Chemicals Ltd.

Preparation of Stem Extract

The stems were washed with distilled water, shade dried and powdered. Powdered drug material was subjected to successive solvent extraction using chloroform, ethyl acetate, methanol, and water respectively. The extracts were filtered, evaporated at 40° and stored at 40°.

Phytochemical Screening

Phytochemical screening was done to determine the presence or absence of secondary metabolites such as tannins, alkaloids, flavonoids, saponins, sterols, and phenolic compounds. This was done according to established procedure [18-21].

Alkaloid Test (Dragendroff's Test)

2 ml plant extract was acidified with few drops of dilute hydrochloric acid. To this acidic medium, 1 ml of Dragendroff's reagent (Potassium bismuth iodide) was added. An orange or reddish brown precipitate produced indicates the presence of alkaloids.

Flavonoid Test (Shinoda Test)

The presence of flavonoids was confirmed by treating the alcoholic plant extract with few fragments of magnesium ribbon and hydrochloric acid. The reaction mixture develops pink or crimson red color, indicating the presence of flavonoids.

Saponin Test (Foam Test)

1 ml of each extract shaken with 10 ml of distilled water and it was agitated in a graduated cylinder for 10 min. The formation of persistent honey-comb like froth indicated the presence of saponins.

Carbohydrate Test (Molish's Test)

A small amount of extract was treated with an alpha-Naphthol solution in alcohol, shakes and adds conc. H₂SO₄ from the side of the test tubes and observed for the formation of the violet ring at the junction of two liquids.

Tannin Test (Lead Acetate Test)

To 2 ml of each extract add a few drops of 10% Lead acetate were added. The appearance of white precipitate indicates the presence of tannins.

Steroids Test

50% H₂SO₄ is added along the sides of the test tube containing a mixture of methanolic HCl and acetic anhydride. If there is any change in color, from green to blue-green (sometimes via red or blue) indicates the presence of terpenoids and steroids.

Phenol Test

When 0.5 ml of FeCl₃ (w/v) solution was added to 2 ml of rest solution, the formation of an intense color indicated the

presence of phenols.

Protein Test (Biuret Test):

To extract add 4% NaOH and a few drops of 1% CuSO₄ solution. The appearance of violet or pink color indicates the presence of Protein.

Anticancer Activity:

The cell lines were grown in RPMI 1640 medium containing 10% fetal bovine serum and 2 mM L-glutamine. For the present screening experiment, cells were inoculated into 96 well microtiter plates in 100 µL at plating densities as shown in the study details above, depending on the doubling time of individual cell lines. After cell inoculation, the microtiter plates were incubated at 37° C, 5% CO₂, 95% air and 100% relative humidity for 24 h before the addition of experimental drugs.

Different extracts were initially solubilized in dimethyl sulfoxide at 100 mg/ml and diluted to 1 mg/ml using water and stored frozen before use. At the time of drug addition, an aliquot of frozen concentrate (1 mg/ml) was thawed and diluted to 100 µg/ml, 200 µg/ml, 400 µg/ml and 800 µg/ml with complete medium containing test article. Aliquots of 10 µl of these different drug dilutions were added to the appropriate microtiter wells already containing 90 µl of the medium, resulting in the required final drug concentrations, i.e. 10 µg/ml, 20 µg/ml, 40 µg/ml, 80 µg/ml.

After compound addition, plates were incubated at standard conditions for 48 h and assay was terminated by the addition of cold TCA. Cells were fixed in-situ by the gentle addition of 50 µl of cold 30 % (w/v) TCA (final concentration, 10% TCA) and incubated for 60 min at 4 °C. The supernatant was discarded; the plates were washed five times with tap water and air dried. Sulforhodamine B (SRB) solution (50 µl) at 0.4% (w/v) in 1% acetic acid was added to each of the wells, and plates were incubated for 20 min at room temperature. After staining, the unbound dye was recovered, and the residual dye was removed by washing five times with 1% acetic acid. The plates were air dried. The bound stain was subsequently eluted with 10 mM trizma base, and the absorbance was read on a plate reader at a wavelength of 540 nm with 690 nm reference wavelength.

Percent growth was calculated on a plate-by-plate basis for test wells relative to control wells. Percent Growth was expressed as the ratio of average absorbance of the test well to the average absorbance of the control wells * 100 [22-25].

Using the six absorbance measurements [time zero (Tz), control growth (C), and test growth in the presence of drug at the four concentration levels (Ti)], the percentage growth was calculated at each of the drug concentration levels. Percentage growth inhibition was calculated as:

$$[\text{Ti}/\text{C}] \times 100\% \text{ (Ref)}$$

Results and Discussion

The result of the preliminary phytochemical analysis is presented in Table 1. The phytochemicals present in the stem of *Lantana camara* L. was found to be alkaloids, saponins, tannins, flavonoid, steroids, and phenolic compounds. The presence of one or more of these secondary metabolites indicated that the anticancer activity is due to these active compounds present in different parts of the plant.

In the present study solvents, extracts of the stem of *Lantana camara* L., namely chloroform, ethyl acetate, methanol, and aqueous extracts were evaluated for anticancer activity on the

cancer cell lines HL-60 (Leukemia) at various concentrations. The activities of the different extracts were compared with the standard drug Adriamycin (Doxorubicin). The result was found that chloroform, ethyl acetate, and methanol extracts were active on Human Leukemia Cell Line HL-60. The GI₅₀ value of chloroform, ethyl acetate, methanol and aqueous extracts for HL-60 cell line was found to be <10 µg/ml except for aqueous extract >80 µg/ml. Doxorubicin served as a positive control which showed GI₅₀ value <10 µg/ml on three cancer cell lines Table 2. The results of in-vitro cancer activity showed that *Lantana camara* L. plant could be used in the treatment of leukemia and the demonstration of anticancer activity against Human Leukemia Cell Line HL-60 is an indication that there is the possibility of sourcing of new anticancer compounds from screened plant leading to the discovery of new compounds.

Table 1: Phytochemical Analysis of Different Extracts of *Lantana Camara* L.

Phytochemicals	Different Extracts of <i>Lantana Camara</i> L.			
	Chloroform	Ethyl Acetate	Methanol	Aqueous
Alkaloids	+	+	+	-
Phenols	+	+	+	-
Tannins	+	+	+	+
Flavonoids	+	+	+	+
Saponins	+	+	+	+
Steroids	+	+	-	-
Carbohydrates	-	-	-	+
Proteins	-	-	-	+

Where, + = Present, - = Absent

Table 2: In-Vitro Anticancer Activity On HL-60 Celllines

Extracts	% Growth Control Drug concentration µg/ml				GI ₅₀
	10	20	40	80	
Chloroform extract	-5.8	-3.5	-10.7	-18.6	<10
Ethyl acetate extract	6.0	27.6	40.3	6.4	<10
Methanolic extract	20.5	37.4	29.5	45.0	<10
Aqueous extract	133.0	91.3	99.3	72.6	>80
ADR	-64.3	-68.7	-72.6	-73.5	<10




Fig 1: Growth Curve: Human Leukemia cancer Cell Line HI-60

Conclusion

In conclusion, *Lantana camara* L. stem can be considered as an important source of natural products that have anticancer potential. Further investigation is needed to be done to find out the active compounds responsible for the anticancer activity and their mechanism of action. Chloroform, ethyl acetate, methanol and aqueous extracts of *Lantana camara* L. stem were tested for their anticancer activity on the cancer cell

lines HL-60 (Leukemia) at various concentrations. The anticancer effect of the extracts on cell inhibition was studied using Sulphorhodamine B assay. Only chloroform, ethyl acetate, and methanolic extracts showed the dose-dependent anticancer effect on leukemia cancer cells. Phytochemical studies of the plant extracts showed the presence of flavonoid, alkaloids, saponins, tannins, and triterpenes. The result revealed that chloroform, ethyl acetate and methanolic extracts of *Lantana camara* L. contains important chemical constituents, which can be further used in the treatment of leukemia. However, the present study of the in-vitro anticancer activity of *Lantana camara* L. forms primary Platform for further phytochemical and pharmacological studies.

References

1. Tyagi N, Sharma GN, Shrivastava B, Saxena P, Kumar N. Medicinal plants: used in Anti-cancer treatment. International Journal of Research and Development in Pharmacy & Life Sciences 2017;6(5):2732-39. doi: 10.13040/IJRDP.L.2278-0238.
2. Mohamed A, El-Awady, Nabil S, Awad, Adel E, E-Tarras. Evaluation of the anticancer activities of pomegranate (*Punica granatum*) and harmful (*Rhazya stricta*) plants grown in Saudi Arabia. International Journal of Current Microbiology and Applied Sciences 2015;4(5):1158-67.
3. Somayeh AV, Meera S. Preclinical evaluation of in-vivo anticancer activity of *Punica grantum* L. and *Ziziphus mauritiana* using Ehrlich ascites carcinoma in Swiss albino mice. European Journal of Pharmaceutical and Medical Research 2016;3(3):402-08.
4. Sumitra C, Krunal N. In-vitro and in-vivo methods for anticancer activity evaluation and some Indian medicinal plants possessing anticancer properties: An Overview. Journal of Pharmacognosy and Phytochemistry 2013;2(2):140-52.
5. Vickers A. Botanical medicines for the treatment of cancer: rationale, Overview of current data, and methodological considerations for phase I and II trials. Cancer Invest 2002;20:1069-79.
6. Rajdeep K, Karan K, Harpreet K. Plants as a source of anticancer agents. Journal of Natural Product and Plant Resources 2011;1:119-24.
7. Krishnamurthy PT, Vardhranjalu A, Wadhawani A, Patel V. Identification and characterization of a potent anticancer fraction from the leaf of *Moringa oleifera* L. Indian Journal of Experimental Biology 2015; 53:98-03.
8. David P: Bengal Plants. Dehradun, India 1981;822-38.
9. Nelson G. The shrubs and woody vines of Florida. Pineapple Press, Inc. Sarasota 1996.
10. Castro O, Barrios M, Chinchilla M, Guerrero O. Chemical and biological evaluation of the effect of plant extracts against *Plasmodium berghei*. Revista De Biologia Tropical 1996;44:361-67.
11. Whistler WA. Tropical ornamentals, a guide. Timber Press, Inc Portland 2000.
12. Xiao P. A pictorial Encyclopedia of Chinese Medical Herbs. Chuokoron-Sha, Inc 1992.
13. Takeda Y, Morimoto Y, Matsumoto T, Ogini C, Hirata E, Takushi A. Iridoid glycosides from the leaves and stem of *Lantana camara* L. Phytochemistry 1995;39(4):829-33.
14. Rimpler H, Timm H. Iridoids and ecdysones from Verbenaceae. V: Iridoids from *Lantana camara* L. Z Naturforsch (C) 1974;29:111-15.

15. Salama OM, Amer MM, Lahloub MF, Spengel S: Repennoside: A new iridiod glucoside from *Lantana camara* L Fruits. *Jou of Pharmaceutical Sci* 1992;8:21221.
16. Anis, I, Ahmed S, Malik A, Yasin A, Choudary MI. Enzyme inhibitory constituents from *Duranta repen*. *Chemical and Pharmaceutical Bulletin* 2000;50:515-518.
17. Subramanian SS, Nair AG. Scutellarein and pectolarigenin from the leaves of *Clerodendron phlomides* and *Lantana camara* L. *Phytochemistry* 1972;50:3095-96.
18. Swamy NT, Rosaiah G, Babu K, Kumar KV. A study on phytochemical composition, GC-MS analysis and antimicrobial potential of methanolic leaf extract of *Alstonia scholaris* (L.) R. Br. *International Journal of Pharmaceutical Sciences & Research* 2019;10(3):747-55. doi:10.13040/IJPSR.0975-8232.
19. Fawole OA, Ndhala AR, Amoo SO, Finnie JF, Van- Staden J. Anti-inflammatory and phytochemical properties of twelve medicinal plants used for treating gastro-intestinal ailments in South Africa. *Journal of Ethnopharmacology* 2009;123:237-43.
20. Ndam LM, Mih AM, Fongod AGN, Tening AS, Tonjock RK, Enang JE. Phytochemical screening of the bioactive compounds in twenty (20) Cameroonian medicinal plants. *International Journal of Current Microbiology and Applied Sciences* 2014;3(12):768-78.
21. Ngbede J, Yakubu RA, Nyam DA. Phytochemical screening for active compounds in *Canarium schweinfurthii* (atile) leaves from Jos North, Plateau State, Nigeria. *Research Journal of Biological Sciences* 2008;3(9):1078-78.
22. Vanicha V, Kanyawim K, Sulforhodamine B. colorimetric assay for cytotoxicity screening. *Nature Protocols* 2006, 1112-16.
23. Skehn P, Storeng R, Scudiero A, Monks J, McMohan D, Vistica D *et al.*: New colorimetric cytotoxicity assay for anticancer drug screening *Journal of the National Cancer Institute* 1990, 1082-07.
24. Chavan R, Khan M, Sathe N, Mankar N. A Review: SrB assay for screening anticancer activity of herbal drugs (*in-vitro*). *International Ayurvedic Medical Journal* 2016;4(2):66-70.
25. Esteban AO, Andrea LK. Sulforhodamine B (SRB) assay in cell culture to investigate cell proliferation. *Bio Protocol* 2016, 6(21). doi:10.21769/Biorotoc.1984.