



E-ISSN: 2278-4136

P-ISSN: 2349-8234

www.phytojournal.com

JPP 2020; 9(4): 317-319

Received: 28-05-2020

Accepted: 30-06-2020

Nidhi Joshi

PG Students, Department of
Biochemistry, College of Basic
Sciences and Humanities,
Govind Ballabh Pant University
of Agriculture & Technology,
Pantnagar (U. S. Nagar),
Uttarakhand, India

Kuldip Chandra Verma

Assistant Professor, College of
Basic Sciences and Humanities,
Department of Biochemistry,
Govind Ballabh Pant University
of Agriculture & Technology,
Pantnagar (U. S. Nagar),
Uttarakhand, India

Corresponding Author:**Kuldip Chandra Verma**

Assistant Professor, College of
Basic Sciences and Humanities,
Department of Biochemistry,
Govind Ballabh Pant University
of Agriculture & Technology,
Pantnagar (U. S. Nagar),
Uttarakhand, India

A review on nutrition value of Amaranth (*Amaranthus caudatus* L.): The crop of future

Nidhi Joshi and Kuldip Chandra Verma

Abstract

From the past 20 years, little progress has been achieved in reducing child malnutrition food insecurity and hunger. To address these problems, increased consumption of leafy vegetables is promoted as sources of both bio-active compounds and micronutrients. Amaranthus is one of the crop, which has been rediscovered as a promising food crop in form of leaf and seed, mainly due to its resistance to drought, heat, diseases and pests, and the high nutritional value of both seeds and leaves. The seeds are small and lent cellular or oval in shape. The grain is high in fibre and low in saturated fats, factors which contribute to its use by the health food market. The oils obtained from amaranth are important because they contain oleic, linoleic and linolenic fatty acids. The lipid fraction of amaranth also contains compounds with antioxidant potentials, such as tocopherols, which act as scavengers of lipid peroxyl radicals. It also contains several classes of flavonoids and anthocyanins, as well as other phenolic constituents. It is high in protein content which contains essential amino acids that are good for human nutrition. It contains a high level of lysine and sulphur-containing amino acids like cysteine. Amaranth has become popular among patients with celiac disease because it does not cause allergic reactions in the intestinal mucosa. Quantification of bio-active compounds of Amaranthus seed oils like squalene, β -sitosterol and α -tocopherol will help enhance its use for human consumption as a nutraceutical. Leaves and succulent stem are good sources of iron, calcium, vitamin A and vitamin C. In view of the enormous nutritional benefits of the crop, there is a definite need for its genetic improvement to develop high yielding varieties with high content of desired quality traits.

Keywords: Amaranth, quality traits, lysine, squalene, bio-active, phenolic

Introduction

In the world child malnutrition, food insecurity and hunger is a big problem. Lack of essential nutrients is spread mainly among children and women and due to which global disease burden of children increases by limiting proper cognitive development, impairing physical development, and increasing susceptibility to infectious diseases. So, In order to overcome these problems increased consumption of seeds and leafy vegetables is very good, as these are considered sources of many vital micronutrients, macronutrients and bio-active compounds. Farming communities and other peoples living in Uttarakhand hills depends heavily on available resources, climate and geographic conditions for food and nutritional security. The climate of Himalayan region having a wide ranges and hence suitable for cultivation of economical and nutritive crops like Grain Amaranthus, Foxtail Millet, Finger Millet, Buck-Wheat, Rice Bran, Barnyard Millet etc. These crops are underutilized but are commercially important. One can get food products with enhanced nutritional value by using them ^[1]. It is mainly due to its resistance to drought, heat, diseases and pests. As Amaranthus exhibit C4 photosynthesis pathway it can tolerate a large variety of unfavourable conditions, including high salinity, alkalinity, or acidity, making it suited for subsistence agriculture. Amaranthus have high nutritional value in both seeds and leaves ^[2]. The genus Amaranthus is a valuable food source of nutrients containing high quality proteins, minerals, vitamins and bioactive compounds such as phenolics. It can be consumed as a staple food, additionally both the grain and leaves of vegetable Amaranthus have also been commercially used for natural dyes, skin lubricants and pharmaceuticals. The grains of Amaranthus are gluten-free, and also contain a balanced essential amino acid profile. Both the grain and leafy vegetable portion of Amaranthus plants have many beneficial health effects. The Amaranthus leaves also show anticancer potential by inhibiting the proliferation of liver, breast and colon cancer cell lines ^[3]. Amaranthus is considered as a native of tropical America and its meaning is immortal in Greek. In the Aztec, Mayan, Incan civilizations it was a staple crop. Now a days, it is cultivated throughout India, China, Nepal, Indonesia, Malaysia, Phillipines; whole of Central America, Mexico; Southern and Eastern Africa. Genus Amaranthus belongs to order Caryophyllales, family Amaranthaceae, sub-family Amaranthoideae. It have 70 species, out of which 40 are

native to the Americas, 17 *Amaranthus* species are of vegetative types, and 3 are grain types (*A. caudatus*, *A. cruentus* and *A. hypochondriacus*)^[4].

Plant morphology

Amaranthus is an annual herbaceous plant which is having height of 1-6 feet, leaves are petioled and alternate having a length of 3-6 inches, vary in colours and rough, ovate, hairy or rhombic with wavy margins. The flowers are small, with greenish or red terminal panicles. Taproot is long, fleshy red or pink. The seeds are small and lent cellular or oval in shape. The colour of seeds may vary from white, red to black^[5]. *Amaranthus* is a non-grass cereal-like (pseudo cereal) dicotyledonous plant. The leaves and grains both of *Amaranthus* are used as staple food in major/minor form^[6]. Pollination in *Amaranthus* is either by wind or by insects, many of the species are monoecious, some are dioecious. The inflorescence can be drooping erect, rope-like, or inconspicuous in the leaf axil. *Amaranthus* has numerous utilities, and based on them it is also known as super grain, miracle grain, and grain of future^[7]. *Amaranthus* leaf and stalk have a wide variety of colours that varies from red to green, and the colour of seed ranges from black to white. Highly pigmented vegetables like *Amaranthus* usually have higher nutritional value as compared to other species of the same plant. The colour of most *Amaranthus* plants is bright red-violet, and betalains are generally considered to be responsible for the maroon pigment^[3].

Nutritional value

In hills, the leaves of *Amaranthus* are used in form of vegetable or saag, and ripen seeds are used in form of very nutritious pseudo cereal with very high amount of protein as compared to the true cereals. High fibre and low saturated fat makes grain *Amaranthus* popular in the health food market. As per the information of national research project, in India, grain *Amaranthus* is recognized as a major crop to produce highly nutritive food and feed. *Amaranthus* is considered to be new millennium crops of nutraceutical value^[8]. The seeds have high protein content having essential amino acids mainly methionine and lysine; it also contains significant levels of squalene, which is an important precursor for all steroids.^[9] Element analysis of *Amaranthus* in mg/100 g (DW) reported that the leaves contained potassium (54.20), sodium (7.43), calcium (44.15), Iron (13.58), Magnesium (231.22), Zinc (3.80) and phosphorus (34.91). The *Amaranthus* leaves have high amount of bio-available iron^[10]. The vitamin content of the leaves in mg/100 g (DW) was -carotene (3.29), thiamine (2.75), pyridoxine (2.33), riboflavin (4.24), niacin (1.54), ascorbic acids (25.40) and tocopherol (0.50). *Amaranthus* contains 17.5 – 38.3 % protein, which consists of 5 % lysine and sulphur-containing amino acids (lacking in cereals and tubers) and many other amino acids are also present in considerable amount^[11]. Seventeen amino acids (isoleucine, leucine, lysine, cysteine, phenylalanine, tyrosine, threonine, methionine, valine, alanine, arginine, glutamic acid, aspartic acid, glycine, histidine, proline and serine) were detected in *Amaranthus*. Lysine content is much higher than maize (3-3.5 times) and wheat (2-2.5 times)^[12]. Biological value of *Amaranthus* protein is 75 and blending of *Amaranthus* with corn flour almost reaches the biological value 100^[13]. *Amaranthus* protein is highly digestible (approx. 90 %). *Amaranthus* protein is gluten free and is therefore suitable for celiac disease patients^[14].

Average oil content of *Amaranthus* varies between 11 to 14%. The oils obtained from *Amaranthus* have a high saponification value (130-190) and the iodine value (100-113). The unsaponifiable matter present is (5-7%) that is of great importance. The major 5 fatty acids present in *Amaranthus* were: Oleic acid (26.5-31%), palmitic acid (14-20%), stearic acid (2-3.5%); linoleic acid (32-41%) and docosahexaenoic acid (DHA) (7-21%)^[15].

Polyphenols and antioxidant properties

Three polyphenols viz. rutin, isoquercitrin, and nicotiflorin were isolated in *Amaranthus*. Rutin was present in higher concentration (10.1 mg/g flour) in seed flour. Eleven phenolic compounds (gallic acid, protocatechuic acid, chlorogenic acid, gentistic acid, 2,4-dihydroxybenzoic acid, ferulic acid, salicylic acid, rutin, ellagic acid, kaempferol-3-rutinoside and quercetin) were recognized in the extracts of different parts of *Amaranthus*. The total phenolic content (TPC) fluctuated from 1.04 to 14.94 mg GAE/g DW; the total flavonoid content (TFC) fluctuated from 0.27 to 11.40 mg CAE/g DW; while the total betalain content (TBC) fluctuated from 0.07 to 20.93 mg/100 g DW. The Total Phenolics content of 8.45 GAE equivalents ($\mu\text{g GAE/mg sample}$) was estimated to be 0.85 mg GAE /100 mg extract^[16]. Antioxidant activity Free phytochemicals content in crude seeds was 2.32 ± 0.12 while after optimization it increased to 11.26 ± 0.20 . Bound phytochemicals content in crude seeds was 3.08 ± 0.12 while after optimization it increased to 10.30 ± 0.21 . Total phytochemicals content in crude seeds was 5.40 ± 0.13 while after optimization it increased to 21.56 ± 0.30 . Antioxidant activity Free phytochemicals content in crude seeds was 0.90 ± 0.06 while after optimization it increased to 7.71 ± 0.13 . Bound phytochemicals content in crude seeds was 1.23 ± 0.09 while after optimization it increased to 4.44 ± 0.11 . Total phytochemicals content in crude seeds was 2.13 ± 0.12 while after optimization it increased to 12.16 ± 0.23 . Total phenolics content in crude seeds was 26.65 ± 1.10 , while after optimization it increased to 247.63 ± 2.20 . Total flavonoids content in crude seeds was 26.01 ± 0.14 while after optimization it increased to 81.39 ± 1.17 ^[17].

Medicinal uses

Amaranthus has astringent, appetizing and diuretic properties and is used in the treatment of diarrhoea, dysentery^[18], inflammation, gonorrhoea and piles^[19]. Whole plant of *Amaranthus* is used as laxative and the root is regarded as highly specific for colic. *Amaranthus* leaves and roots are traditionally boiled and given to children as laxative. *Amaranthus* is also used traditionally as diuretic, anti-snake venom, antipyretic, antidiabetic, antigonorrhoea, and antileprotic. The plant sap is used as an eye-wash to treat ophthalmia and convulsions in children. The seeds of *Amaranthus* or oil are beneficial for those with hypertension and cardiovascular diseases; it also reduces high blood pressure and cholesterol levels, and also helps in subsiding antioxidant level in body and immune system. The leaves of *Amaranthus* are recommended well for young children, lactating mothers and for patients with constipation, haemorrhage, fever, anemia or kidney complaints. The water of softened plants is used as to treat pains in the limbs. The ash of the stems is used in wound dressing. Heated leaves can be used in tumours. *Amaranthus* can be used externally to treat inflammations, and internally to treat diuretic problems. The seed of *Amaranthus* is used as a poultice for broken bones. It is used internally in the treatment of internal

bleeding and excessive menstruation. Root can be used in treatment of gonorrhoea and is also applied as an emmenagogue and antipyretic. Crushed leaves are also considered a good emollient. It is also used for the treatment of snake-bites [2].

Conclusion

Amaranthus is a source of various phytochemicals (rutin, isoquercitrin, nicotiflorins etc.). It having tremendous used in our daily food and industries. These phytochemicals also having various uses in medical field. *Amaranthus* has therefore a great potential, because it is a sustainable crop that could help in improving household food security and income of farmers. In arid regions where commercial crops such as maize, beans or rice cannot be grown, *Amaranthus* cultivation could contribute to the first Millennium Development Goal of reducing hunger and poverty. To improve breeding of *Amaranthus* the research should be strengthened and that requires collection and diffusion of knowledge of many local communities on the value of neglected *Amaranthus* species. Identification and preservation of germplasm is very important for maintaining genetic diversity and studying genetic material to improve the nutritional quality. Inter-specific and intra-specific hybridisation should be done to improve seed yield, macro and micro-nutrients of cultivated *Amaranthus* including vitamin A, iron and zinc contents. There should be an effective collaboration between farmers and scientists for proper utilization of *Amaranthus* crop.

References

1. Mlakar SG, Turinek M, Jakop M, Bavec M, Bavec F. Nutrition value and use of grain amaranth: potential future application in bread making. *Agricultura*. 2009; 6(2):45-53.
2. Achigan-Dako EG, Sogbohossou OE, Maundu P. Current knowledge on *Amaranthus* spp.: research avenues for improved nutritional value and yield in leafy amaranths in sub-Saharan Africa. *Euphytica*. 2014; 197(3):303-317.
3. Hongyan L, Deng Z, Liu R, Zhu H, Draves J, Marcone M et al, Tsao R. Characterization of phenolics, betacyanins and antioxidant activities of the seed, leaf, sprout, flower and stalk extracts of three *Amaranthus* species. *Journal of Food Composition and Analysis*. 2015; 37:75-81.
4. Ebert AW, Wu TH, Yang RY. Amaranth sprouts and microgreens—a homestead vegetable production option to enhance food and nutrition security in the rural-urban continuum. In Proceedings of the Regional Symposium on Sustaining Small-Scale Vegetable Production and Marketing Systems for Food and Nutrition Security (SEAVEG 2014), Bangkok, Thailand, 2014, 25-27.
5. Akubugwo IE, Obasi NA, Chinyere GC, Ugbogu AE. Nutritional and chemical value of *Amaranthus hybridus* L. leaves from Afikpo, Nigeria. *African Journal of Biotechnology*. 2007; 6(24):2833-2839
6. Tang Y, Li X, Chen PX, Zhang B, Liu R, Hernandez M, Tsao R. Assessing the fatty acid, carotenoid, and tocopherol compositions of amaranth and quinoa seeds grown in Ontario and their overall contribution to nutritional quality. *Journal of agricultural and food chemistry*. 2016; 64(5):1103-1110.
7. Ofitserov EN. Amaranth: Perspective raw material for food processing and pharmaceutical industry. *Chemistry and Computational Simulation*. Butlerov Communications. 2001; 2(5):1-4.
8. Rastogi A, Shukla S. Amaranth: a new millennium crop of nutraceutical values. Critical reviews in food science and nutrition. 2013; 53(2):109-125.
9. De la Rosa AP, de León-Rodríguez A, Laursen B, Fomsgaard IS. Influence of the growing conditions on the flavonoids and phenolic acids accumulation in amaranth (*Amaranthus hypochondriacus* L.) leaves. *Terra Latinoamericana*. 2019; 37(4):449-57.
10. Rangarajan A, Kelly JF. Further studies exploring iron nutritional quality from *Amaranthus* species. *Horticulture Science*. 1994; 29(5):458-460.
11. Kumar BA, Lakshman K, Jayaveera KN, Khan S, Manoj B, Swamy VBN. Evaluation of the antioxidant activity of *Amaranthus spinosus* Linn. by non-enzymatic haemoglycosylation. *Sains Malaysiana*. 2010; 39(3):413-415.
12. Svirskis A. Investigation of amaranth cultivation and utilization in Lithuania. *Agronomy research*. 2003; 1(2):253-264.
13. Oboh BO, Aghalokpe IO. Seed Protein Characterisation of Some Selected Cultivated *Amaranthus* Accessions. *Journal of Science Research and Development*. 2002; 7:9-16.
14. Alvarez-Jubete L, Arendt EK, Gallagher E. Nutritive value of pseudocereals and their increasing use as functional gluten-free ingredients. *Trends in Food Science & Technology*. 2010; 21(2):106-113.
15. Dhellot JR, Matouba E, Maloumbi MG, Nzikou JM, Ngoma DS, Linder M et al. Extraction, chemical composition and nutritional characterization of vegetable oils: Case of *Amaranthus hybridus* (var 1 and 2) of Congo Brazzaville. *African Journal of Biotechnology* 2006; 5(11):1095-1101
16. Bhat A, Satpathy G, Gupta RK. Evaluation of Nutraceutical properties of *Amaranthus hypochondriacus* L. grains and formulation of value added cookies. *Journal of Pharmacognosy and Phytochemistry* 2015; 3(5):51-54
17. Perales-Sánchez JX, Reyes-Moreno C, Gómez-Favela MA, Milán-Carrillo J, Cuevas-Rodríguez EO, Valdez-Ortiz A et al. Increasing the antioxidant activity, total phenolic and flavonoid contents by optimizing the germination conditions of amaranth seeds. *Plant foods for human nutrition*. 2014; 69(3):196-202.
18. Sarkar R, Nandan CK, Mandal S, Patra P, Das D, Islam SS. Structural characterization of a heteropolysaccharide isolated from hot water extract of the stems of *Amaranthus tricolor* Linn. (*Amaranthus gangeticus* L.). *Carbohydrate research*. 2009; 344(17):2412-2416.
19. Ishtiaq S, Ahmad M, Hanif U, Akbar S, Kamran SH. Phytochemical and in vitro antioxidant evaluation of different fractions of *Amaranthus graecizans* subsp. *silvestris* (Vill.) Brenan. *Asian Pacific journal of tropical medicine*. 2014; 7:342-347.