



E-ISSN: 2278-4136

P-ISSN: 2349-8234

www.phytojournal.com

JPP 2021; Sp 10(1): 143-145

Received: 07-11-2020

Accepted: 09-12-2020

Heena J Madavi,
M.Sc. Student, Department of
Animal Husbandry and Dairy
Science, COA, Latur,
Maharashtra, India

Akshay P Deshmukh
Assistant Professor, Department
of Animal Husbandry and Dairy
Science, BCOA, Selu,
Maharashtra, India

AG Sarnaik
Assistant Professor, Department
of Animal Husbandry and Dairy
Science, RGDCOA, Tivsa,
Maharashtra, India

PD Sarnaik
M.Sc. Student, Department of
Animal Husbandry and Dairy
Science, COA, Latur,
Maharashtra, India

RA Patil
Assistant Professor, Department
of Animal Husbandry and Dairy
Science, VNMKVCOA,
Parbhani, Maharashtra, India

Corresponding Author:
Akshay P Deshmukh
Assistant Professor, Department
of Animal Husbandry and Dairy
Science, BCOA, Selu,
Maharashtra, India

Combined effect of drumstick (*Moringa olifera*) and lemongrass (*Cymbopogon citratus*) on growth performance of broilers

Heena J Madavi, Akshay P Deshmukh, AG Sarnaik, PD Sarnaik and RA Patil

Abstract

An experiment was conducted at poultry unit in college of Agriculture Latur, to determine the combined effect of Drumstick (*Moringa olifera*) and Lemongrass (*Cymbopogon citratus*) powder on growth performance and economical of feeding. Eighty day-old chicks were divided into four dietary treatments i.e., T₁: Basal diet feeding, T₂: Basal diet+ 0.5% Drumstick powder, T₃: Basal diet+0.5% Lemongrass powder and T₄: Basal diet +0.5% Drumstick+ 0.5% Lemongrass powder of combination with four replication and five birds have each replication. Result of experiment showed significant difference (P<0.05) in feed intake, live weight, body weight gain and feed conversion ratio between treatments. The study concluded that addition of drumstick powder at the 0.5 per cent and 1 per cent mixture of Drumstick and lemongrass in broiler diet improved growth performance of broiler and economical and profitable for broiler production.

Keywords: Broilers, economics, FCR, natural feed additives, weight gain

Introduction

Antibiotics have been used widely to prevent infections and poultry diseases and for the improvement of meat and egg production. However, use of antibiotics is restricted due to drug resistance in bacteria, drug residue in carcass and also alteration of natural gut micro flora. Recently many countries tend to minimize or prohibit the use of antibiotics because of their deleterious side effects on both animals and human. Consequently, the use of natural promoters such as probiotics, prebiotics, symbiotics, enzymes, toxic binders, organic acids, oligosaccharides, phytogenics, and other feed additives, to enhance the growth and performance of broiler chickens have been advocated compounds. The lemongrass metabolites as oil (LGO) is considered as a viable alternative to antibiotics for the broiler and have been studied as an alternative for microbial and growth promoting abilities in the poultry and that resulted minimized feed expense in the production chain Drumstick (*Moringa olifera*): Moringa is a potential plant that could be used to enhance immune response and to improve intestinal health of broiler chicken. The leaves of *Moringa* has high protein content which is between 20-33% on a dry weight basis, the protein is of high quality having significant qualities of all the essential amino acid

Materials and Methods

The present research work was undertaken to study the combined effect of drumstick and lemongrass supplementation on growth performance of broilers. The present study was carried out in the Department of Animal Husbandry and Dairy Science, College of Agriculture, Latur, VNMKV, Parbhani, Maharashtra state. Day old eighty, commercial straight run broiler chicks (Vencobb-430) strains were obtained from Huma hatcheries Latur (Maharashtra). Weighted (46.71±0.02g) and randomly allocated into four treatments groups with four replications and each replication have with five birds.

Experimental design

The concentration of the administrated Additives in four experimental diets were as follows: T₁ - Standard broiler ration without supplement (control), T₂ - 99.50 part standard broiler ration + 0.5 part drumstick leaf meal, T₃ - 99.50 part standard broiler ration + 0.5 part lemongrass leaf meal and T₄ - 99.00 part standard broiler ration + 0.5 part mixture of drumstick leaf meal + 0.5part of lemongrass leaf meal. The herbal growth promoter (Drumstick and Lemongrass) was obtained from local market of Latur Dist, Latur, Maharashtra.

Weekly growth of chicks and daily feed consumption in each group were recorded up to six-week period. Feed conversion efficiency (FCR) was calculated as the ratio between unit feed consumed to unit body weight gain.

The data generated from the experiment were subjected to statistical analysis using randomized block.

Birds Management

Day old eighty, commercial straight run broiler chicks (Vencobb-430) strains were obtained from Huma hatcheries Latur (Maharashtra). Weighted (46.71+0.02g) and randomly allocated into four treatments groups with four replications and each replication have with five birds.

The concentration of the administrated Additives in four experimental diets were as follows: T₁ - Standard broiler ration without supplement (control), T₂ - 99.50 part standard broiler ration + 0.5 part drumstick leaf meal, T₃ - 99.50 part standard broiler ration + 0.5 part lemongrass leaf meal and T₄ - 99.00 part standard broiler ration + 0.5 part mixture of drumstick leaf meal + 0.5part of lemongrass leaf meal.

The herbal growth promoter (Drumstick and Lemongrass) was obtained from local market of Latur Dist, Latur, Maharashtra. Birds were vaccinated routinely against infectious new castle Disease (lasota) 7thday vaccine by intraocular route, 14thday Gambaro vaccine by intraocular route and 21thday Booster dose of Lasota vaccine through intraocular route. The experimental diets and drinking water were provided ad labium The present research work was undertaken to study the combined effect of drumstick and lemongrass supplementation on growth performance of broilers.

The present study was carried out in the Department of Animal Husbandry and Dairy Science, College of Agriculture, Latur, VNMKV, Parbhani, Maharashtra state. throughout the experimental period of six week.

All chickens were maintained at uniform temperature and lighting control system during the whole period of study. Weekly growth of chicks and daily feed consumption in each group were recorded up to six week period. Feed conversion efficiency (FCR) was calculated as the ratio between unit feed consumed to unit body weight gain.

The data generated from the experiment were subjected to statistical analysis using randomized block.

Proximate Analysis

Proximate analysis of drumstick and lemongrass leaf meal were performed for the components of dry matter, crude protein, ether extract, crude fiber and ash, Proximate analysis of drumstick and lemongrass leaf meal were performed for the components of dry matter, crude protein, ether extract, crude fiber and ash, according to In Table 2 showed chemical composition of natural feed additives.

Body weight gain of broiler chicks (g/bird/week) as affected by addition of drumstick and lemongrass leaf meal.

Items	0%	0.5%	0.5%	Drumstick + lemongrass			
					1%	SE	CD at 5%.
1 st week	134.79	143.60	137.79	142.06	6.35	N.S.	
2 nd week	259.80	269.80	262.25	267.50	11.03	N.S.	
3 rd week	399.20	420.00	424.00	422.00	15.69	N.S.	
4 th week	499.75	540.00	502.00	537.00	22.66	N.S.	
5 th week	439.75	590.00	500.00	585.00	30.5	N.S.	
6 th week	628.00						
Total	2361.29 ^a	2583.40 ^b	2430.04 ^{ab}	2568.56 ^b	54.51	159.61*	

NS = no significant difference ($P>0.05$) *Significant different ($P<0.05$)

Diets formulated to meet the nutrient requirements of broilers according to the recommendations of Ross Broiler manual.

Results and Discussion

To asses combined effect of drumstick and lemongrass, weight gain of Eighty, day old, broiler improved strain (Vencobb 430) were distributed in to four equal groups with four replication of 20 chicks.

The experimental broiler chicks were reared as per standard on deep litter system in well ventilated shed from 0-6 weeks.

Per cent chemical composition of experimental broiler ration on dry matter basis

Sr. No.	Nutrients	Per cent in ration		
		Pre-starter	Starter	Finisher
1.	Crude protein	23.35	21.64	20.20
2.	Crude fiber	3.92	3.96	3.99
3.	Ether extract	4.95	5.08	5.12
4.	Total ash	6.2	6.1	5.93
5.	Acid insoluble ash	1.57	1.60	1.62
6.	Nitrogen free extract	61.58	63.22	64.76
7.	Metabolizable energy (kcal/kg)	2982.5	3065.7	3198.8
8.	E/P ratio	127.73:1	141.66:1	158.35:1

Weight gain

The mean average cumulative body weights of broilers in the treatment groups T₂ was significantly ($P<0.05$) higher as compared to those in T₁, T₂, were as at par with T₄ group. It could be seen that highest cumulative body weight of 2630.25 g obtained in T₂ group broiler receiving 0.5 per cent drumstick leaf meal powder followed by 2615g with 1.00 per cent mixture of drumstick and lemongrass in T₄ and 2476 g with 0.5 per cent lemongrass leaf meal in T₃ and lowest cumulative body weight i.e. 2428 g in T₁ control at the end of 6th week.

The result obtained in those study corresponds with the results of Jamel *et al.*, (2013) also observed that in four dietary treatments consisted of a control (basal diet), basal diet + 0.5% garlic powder , basal diet +0.5% black seed and basal diet +0.5% plant premix and advised to supplement broiler feed with 0.5% for growth performance. Mawahib *et al.* (2016) showed that supplementing broiler diets with 0.2% mixture of garlic and ginger significantly improed live body weight (g/bird) Doaa and Ghada (2015), Results in body weight gain are given in Table 1.

The data showing after the 6th week it was seen from the observations that the average gain in body weight of bird among treatment groups T₂ was significantly superior ($P<0.05$) over (control) group, T₄ whereas at par with treatment T₃ group. The significant ($P<0.05$) positive effect of ginger on body weight gain in weeks 2, 4 and 6

Conclusion

The addition of 0.5 per cent of drumstick 1per mixture of lemon grass and drumstick in broiler ration was beneficial in improving live weight and weight gain in t² and t⁴

References

- 1 Bhalsing GR. Effect of Spirulina Supplementation on Growth Performance of Giriraja Chiken. MSc. Thesis, submitted to the VNMKV 2016. Parbhani-431402.
- 2 Bamidele O, Adejumo IO. Effect of Garlic (*Allium sativum*) and Ginger (*Zingiber officinale* Roscoe) Mixture on Performance Characteristics and Cholestrol Profile of Growing Pullets. International journal of poultry science 2012;11(3):217-220
- 3 Chioma Gibson O, Akinsoyinu Akintunde O, Tayo Grace O, Akinboye Olufusnso E, Afodu Osagie J, Nadubuisi-Ogbonna Lois C et al. Serum Biochemistry and Sensory Evaluation of Broiler Chicken Fed *Cymbopogon citratus* Leaf meal. World Journal of Agricltural Research 2017;5(6):305-309.
- 4 Gadzirayi CT, Masamha B, Mupangwa JF, Washaya S. Performance of Broiler Chickens Fed on Mature *Moringa oleifera* Leaf Meal as a Protein Supplement to Soybean Meal. International Journal of Poultry Science 2012;11(1):5-10.
- 5 Gaikwad BS. Effect of Fenugrek (*Trigonella foenum-graceum* L.) Seed Powder as a Natural Feed Additive on Growth Performance of Broilers. MSc. Thesis, Submitted to the VNMKV 2018. Parbhani-431402
- 6 Gujarwar SV. Effect of Ginner (*Zingiber officinale*) As a Natural Feed Additive on Growth Performance of Broilers. MSc. Thesis, Submitted To the VNMKV Parbhani, [MS] 2018.