



E-ISSN: 2278-4136

P-ISSN: 2349-8234

[www.phytojournal.com](http://www.phytojournal.com)

JPP 2022; 11(1): 189-191

Received: 16-11-2021

Accepted: 18-12-2021

**Lalit Kumar**

Department of Horticulture,  
Babasaheb Bhimrao Ambedkar  
University, Vidya-Vihar, Rae  
Bareilly Road, Lucknow,  
Uttar Pradesh, India

**Sanjay Kumar**

Department of Horticulture,  
Babasaheb Bhimrao Ambedkar  
University, Vidya-Vihar, Rae  
Bareilly Road, Lucknow,  
Uttar Pradesh, India

**Dr. R. S. Verma**

Assistant Professor,  
School of Agricultural Sciences &  
Technology (SAS&T), Lucknow,  
Uttar Pradesh, India

**Shatrunjay Yadav**

Department of Horticulture,  
Babasaheb Bhimrao Ambedkar  
University, Vidya-Vihar, Rae  
Bareilly Road, Lucknow,  
Uttar Pradesh, India

**Vipnesh Singh**

Department of Horticulture,  
Babasaheb Bhimrao Ambedkar  
University, Vidya-Vihar, Rae  
Bareilly Road, Lucknow,  
Uttar Pradesh, India

**Corresponding Author:****Lalit Kumar**

Department of Horticulture,  
Babasaheb Bhimrao Ambedkar  
University, Vidya-Vihar, Rae  
Bareilly Road, Lucknow,  
Uttar Pradesh, India

## Effect of plant growth regulators and mulches on growth and yield of strawberry (*Fragaria X ananassa* Duch.) Cv. Chandler

Lalit Kumar, Sanjay Kumar, Dr. RS Verma, Shatrunjay Yadav and Vipnesh Singh

DOI: <https://doi.org/10.22271/phyto.2022.v11.i1c.14340>

**Abstract**

A field experiment was undertaken at B.B.A. University, Lucknow (U.P.) during the year 2018-19 and 2019-20 to study the effect of plant growth regulators and mulches on growth and yield of strawberry (*Fragaria x ananassa* Duch.) cv. Chandler. The experiment was laid out in randomized block design with seventeen treatments and three replications. The results revealed that the different plant growth regulators and mulches affect on growth and yield of strawberry. Maximum plant height (24.4cm), number of leaves per plant (28.5), length of leaves (9.4cm) and length of the petiole (12.4cm), number of flower per plant (30.1), number of fruit per plant (24.1) and fruit weight per berry (13.6g) were recorded with GA3 75ppm + black polyethylene. On the other hand, minimum values of these parameters were recorded in treatment control. The minimum days taken to produce the first flower (31.3days), days taken to 50% flowering (43.0days) were recorded with NAA 20ppm +black polyethylene, and the maximum days recorded with treatment control.

**Keywords:** strawberry, PGRs, mulching, growth, yield characters

**Introduction**

Strawberry (*Fragaria x ananassa* Duch.) is an important fruit crop of the family Rosaceae. The attractive red colour, pleasant aroma, and high nutritional value in terms of vitamins (A and C) and minerals (Fe and K) make this fruit highly prized in global markets. Besides the fresh consumption, strawberry fruits are also being used to prepare jam and jellies due to the presence of a high amount of pectin. Strawberry fruits have a fair amount of natural antioxidants that are useful for relieving oxidative stresses (Sharma and Thakur, 2008). Consumption strawberry berry helps in the prevention of various types of cancers and heart-related diseases. This fruit is also reported to be beneficial in the reduction of inflammation and obesity-related disorders (Arfin *et al.*, 2016) <sup>[1]</sup>. For these reasons the demand for strawberries in the markets is increasing gradually and the growers are also encouraged to cultivate this crop to gain higher income. Scientific evidence has also suggested that the strawberry plant responded well to growth regulator application (Jain and Dashora, 2011). Among the plant growth regulators, naphthalene acetic acid (NAA) and gibberellins (GA3) have been widely tested in the modern agricultural system due to their suitability of application at a cheaper rate. The role of these plant growth regulators has been investigated in several fruits by (Bist *et al.*, 2018) <sup>[2]</sup>.

Strawberry is a low surface creeping herb and, hence, mulching plays a very important role in soil moisture conservation, weed control, and regulation of soil hydrothermal regimes. In addition, mulching also improves the vegetative growth of the plant and the quality of strawberry. Better vegetative growth is important for healthy and vigorous runner production in strawberry where the production of quality planting material is a great concern. However, there are very few reports on the effect of mulches on vegetative growth and runner production of strawberries. Therefore, an investigation was conducted to study the effect of various growth regulators and mulches on the vegetative growth of strawberry.

**Materials and Methods**

The experiment was carried out at the Horticulture Research Farm Babasaheb Bhimrao Ambedkar University, Lucknow (U.P.) during the year 2018-19 and 2019-20. The experimental site is situated at 26°50' N, latitude, 80°52' E, longitude, and altitude of 111

meters above mean sea level (MSL). The experiment was laid out in a randomized block design with seventeen treatment combinations i.e., T<sub>1</sub> Control, T<sub>2</sub>GA<sub>3</sub> 50ppm + black polyethylene, T<sub>3</sub>GA<sub>3</sub> 75ppm + black polyethylene, T<sub>4</sub>GA<sub>3</sub> 50ppm + transparent polyethylene, T<sub>5</sub>GA<sub>3</sub> 75ppm + transparent polyethylene, T<sub>6</sub>GA<sub>3</sub> 50ppm + paddy straw, T<sub>7</sub>GA<sub>3</sub> 75ppm + paddy straw, T<sub>8</sub>GA<sub>3</sub> 50ppm + rice husk, T<sub>9</sub>GA<sub>3</sub> 75ppm + rice husk, T<sub>10</sub>NAA 20ppm +black polyethylene, T<sub>11</sub>NAA 40ppm +black polyethylene, T<sub>12</sub>NAA 20ppm + transparent polyethylene, T<sub>13</sub> NAA 40ppm + transparent polyethylene, T<sub>14</sub>NAA 20ppm + paddy straw, T<sub>15</sub>NAA 40ppm + paddy straw, T<sub>16</sub>NAA 20ppm + rice husk and T<sub>17</sub>NAA 40ppm + rice husk. Each replicated three times sixteen plants per plot. The runners of strawberry were planted at a spacing of 30 x 30 cm. The uniform dose of FYM 20 t/ha and N (80 kg/ha) as Urea, P<sub>2</sub>O<sub>5</sub> (40 kg/ha) as SSP, and K<sub>2</sub>O (50 kg/ha) as M.O.P. were applied at the time of field preparation. Black and transparent polyethylene of 200 gauge paddy straw and sawdust were spread in plots. The plant growth regulators were applied on the 30<sup>th</sup> and 45<sup>th</sup> days after planting on strawberry. The required quantity of gibberellic acid and naphthalene acetic acid was measured by measuring cylinder and dissolved in a small quantity of absolute ethyl alcohol and NH<sub>4</sub>OH, respectively and then the final volume was made up to one liter with distilled water. The stock solution of the chemicals was diluted with tap water for preparing the prepared foliar spray required to strengthen sprayed on the upper surface of the plant thoroughly with the help of a hand sprayer. The observations were recorded on various parameters like plant height, number of leaves/plant, length of leaf, length of petiole, days taken to first flower, days took to 50% flower, number of flowers/plant, number of fruits per plant, and fruit weight per berry.

### Result and Discussion

During the course of the investigation, it has been observed was a significant increase in plant height with the application of different types of mulching and plant growth regulators in combination, GA<sub>3</sub> 75ppm + black polythene was the most effective combination in retaining maximum (24.4 cm) plant height followed by (24.3 cm), with GA<sub>3</sub> 50ppm + black polyethylene while the minimum plant height (14.8 cm) was recorded with treatment control during two years. The possible reason for increased plant height might be due to the application of mulching and plant growth regulators treatments which led to a congenial environment in the root zone due to lower weed population, optimum soil moisture level, increased availability of nutrients, and favorable soil temperature, and regulated growth of strawberry plant by causing cell elongation and a corresponding increase in length

of petiole by application of GA<sub>3</sub>. These results are in consonance with that Tiwari *et al.* (2017) [12]

A maximum number of leaves per plant (28.0), length of leaves (9.4 cm), and length of the petiole (12.4 cm) were observed in (GA<sub>3</sub> 75ppm + black polyethylene) followed by (26.4, 9.2, and 12.1 cm) with GA<sub>3</sub> 50ppm + black polyethylene. The possible reason maximum number of leaves the ability of gibberellins to stimulate the process of cell division and expansion in epidermal and parenchyma cells has been well documented (Bist *et al.*, 2018) [2]. Such activities in the meristematic tissue of leaf primordial in GA<sub>3</sub> treated plants might be higher and perhaps a greater number of leaves with broader leaf lamina and petiole of longer length. A higher concentration of GA<sub>3</sub> increases the above mechanisms many folds. Earlier findings also suggested that exogenous application of GA<sub>3</sub> induced a higher number of leaves (Kaur *et al.*, 2017) [7]. However, a very high concentration of GA<sub>3</sub> (125 mg<sup>-1</sup>) resulted in slightly stunted growth in strawberry plants. Since, application of GA<sub>3</sub> at high concentration is reported to have an inhibitory action in plants (Hedden and Sponsel, 2015) [4].

The minimum days taken to first flower (31.3 days) and (43.0 days) in 50% flowering the present investigation were recorded with the application of NAA 20ppm +black polyethylene possible reasons for minimum days taken to first flowering and 50% flowering because of generally auxin and particularly NAA induces flowering by stimulating florigen which moved from petiole to growing tip and converts vegetative bud to flowering bud and fruit set refers to the change in the ovary leading to the development of the fruit. These changes are usually induced after pollination and fertilization which is triggered by NAA. The results are congruent with Kumar *et al.*, 2011 [5].

The maximum number of flowers (30.1) and a number of fruits per plant (24.1) were found with the application of GA<sub>3</sub> 75ppm + Black polyethylene. The number of flowers and fruits per plant because of hormone application accelerated the development of differentiated inflorescence and stimulated flowering. Mulching provided optimum soil moisture and temperature due to the application of GA<sub>3</sub> and black polyethylene mulch as reported by Kumar *et al.*, (2012) [6] in strawberry. Application of GA<sub>3</sub> 75ppm + Black polyethylene recorded maximum fruit weight per berry (13.6 g) with respect to other treatments. This might be due to fact that such treatment-induced good vegetative growth and flower bunch hence initiated a higher number of flowers and percent berry set. This could also be attributed to the improvement in the water, which might increase the photosynthetic rate causing maximum fruit weight. These observations are supported by the findings of Nor *et al.* (2014) [9].

**Table 1:** Effect of plant growth regulators and mulches on growth and yield of strawberry cv. Chandler (mean of 2 years)

Treatment Combinations	Plant height (cm)	Number of leaves per plant	Length of the leaf (cm)	Length of the petiole (cm)	Days took to first flower	Days took to 50% flowering	Number of flowers per plant	Number of fruits per plant	Fruit weight per berry (g)
T <sub>1</sub> Control	18.8	20.5	7.1	9.4	38.3	49.0	12.1	8.4	10.0
T <sub>2</sub> GA <sub>3</sub> 50ppm + Black polyethylene	24.3	26.4	9.2	12.1	34.8	45.1	28.5	22.9	13.4
T <sub>3</sub> GA <sub>3</sub> 75ppm + Black polyethylene	24.4	28.0	9.4	12.4	35.1	45.4	30.1	24.1	13.6
T <sub>4</sub> GA <sub>3</sub> 50ppm + Transparent polyethylene	24.1	25.5	9.0	11.5	35.4	45.9	26.5	22.3	12.6
T <sub>5</sub> GA <sub>3</sub> 75ppm + Transparent	24.2	25.8	9.2	11.7	35.8	45.9	28.0	22.6	13.0

polyethylene									
T <sub>6</sub> GA <sub>3</sub> 50ppm + Paddy straw	23.4	25.2	8.5	10.9	36.1	46.6	25.3	18.5	12.1
T <sub>7</sub> GA <sub>3</sub> 75ppm + Paddy straw	23.9	25.2	8.8	11.1	36.4	46.9	25.0	19.3	12.3
T <sub>8</sub> GA <sub>3</sub> 50ppm + Rice husk	22.4	24.5	8.3	10.6	36.8	46.9	23.0	17.8	11.7
T <sub>9</sub> GA <sub>3</sub> 75ppm + Rice husk	23.1	25.0	8.5	10.7	37.3	47.2	24.0	18.4	11.8
T <sub>10</sub> NAA 20ppm +Black polyethylene	21.7	24.3	8.3	10.5	31.3	43.0	22.6	17.3	11.6
T <sub>11</sub> NAA 40ppm +Black polyethylene	21.6	24.1	8.1	10.5	31.8	43.3	22.3	16.9	11.4
T <sub>12</sub> NAA 20ppm + Transparent polyethylene	21.6	23.7	7.7	10.3	32.5	43.5	21.8	15.6	11.2
T <sub>13</sub> NAA 40ppm + Transparent polyethylene	21.1	22.6	7.6	10.2	32.8	43.8	21.1	14.9	10.9
T <sub>14</sub> NAA 20ppm + Paddy straw	20.9	22.4	7.5	10.1	33.3	44.0	19.1	12.1	10.9
T <sub>15</sub> NAA 40ppm + Paddy straw	20.3	22.1	7.4	9.9	33.8	44.2	18.5	11.8	10.7
T <sub>16</sub> NAA 20ppm + Rice husk	19.8	21.9	7.3	9.7	34.1	44.7	16.6	11.4	10.4
T <sub>17</sub> NAA 40ppm + Rice husk	19.1	20.9	7.3	9.6	34.6	45.0	15.8	10.4	10.2
CD (P=0.05)	1.30	1.72	0.64	1.02	1.43	1.59	1.59	1.45	1.26

## References

- Afrin S, Gasparrini M, Forbes-Hernandez TY, Reboredo-Rodriguez P, Mezzetti B, Varela-Lopez *et al.* Promising health benefits of the strawberry: a focus on clinical studies. *Journal of Agricultural and Food Chemistry*. 2016;64(22):4435-4449.
- Bist TS, Rawat L, Chakra Borty B, Yadav V. A recent advance in the use of plant growth regulators (PGRs) in fruit crops - A review. *International Journal of Current Microbiology and Applied Sciences*. 2018;7(5):1307-1336.
- Dubey V, Meena ML, Tirpathi VK. Effect of plant bio-regulators and micro-nutrient on vegetative growth, yield, and quality of strawberry cv. Chandler. *Progressive Research-An International Journal* 2017;12(3):330-332.
- Hedden P, Sponsel V. A century of gibberellins research. *Journal of Plant Growth Regulation*. 2015;34(4):740-760.
- Kumar R, Saravanan S, Bakshi P, Srivastava JN. Influence of plant growth regulators on growth, yield, and quality of strawberry (*Fragaria x ananassa* Duch) cv. Sweet Charlie. *Progressive Horticulture*. 2011;43(2):264-267.
- Kumar, Bakshi M, Singh DB. Influence of plant growth regulators on growth, yield, and quality of strawberry under U.P. subtropics. *Asian Journal of Horticulture*; 2012;7(2):434-436.
- Kaur P, Kaur A. Effect of various mulches on the growth and yield of strawberry cv. Chandler under sub-tropical conditions of Punjab. *International Journal of Recent Trends in Science and Technology*. 2017;25(1):21-25.
- Khunte SD, Kumar A, Ansari N, Saravanan S. Influence of PGRs and poultry manure on physico-chemical parameters of strawberry (*Fragaria x ananassa* Duch.) Chandler. *Int. J Curr. Microbiol. App. Sci*. 2019;8(12):108-117
- Nor SS, Razifzh MR, Mamat AS, Adzemi MA. Application of gibberellic acid in stem cutting of dragon fruit (*Hylocereus polyrhizus*) Effect on fruit quality and yield at harvest. *Journal of Biology Agriculture and Healthcare*. 2014;4(21):51-55.
- Palei S, Das AK, Sahoo AK, Dash AK, Swain S. Influence of plant growth regulators on strawberry (*Fragaria x ananassa* Duch.) cv. Chandler under Odisha conditions. *International Journal of Recent Scientific Research*. 2016;7(4):9945-9948.
- Tariq S, Bano A, Qureshi KM. Response of strawberry (*Fragaria x ananassa*) cv. Chandler to different mulching materials," *Science, Technology and Development*, vol. 2016;35(3):117-122.
- Tiwari AK, Saravanan S, Lall D. Influence of different plant growth regulators on vegetative growth and Physico-chemical properties of strawberry (*Fragaria x ananassa*) cv. Chandler. *Plant Archive*. 2017;17(1):367-370.
- Uddin AJ, Hossan MJ, Islam MS, Ahsan MK, Maharajah Strawberry growth and yield responses gibberellic acid concentrations. *Journal of Experimental Biosciences*. 2012;3:51-56.
- Vishal VC, Thippesha D, Chethana K, Maheshgowda BM, Veeresha BG, Basavraj AK. Effect of various growth regulators on vegetative parameters of strawberry (*Fragaria x ananassa* Duch.) cv. Sujatha. *Research Journal Chemical and Environmental Sciences* 2016;4(4):68-71.