



E-ISSN: 2278-4136  
P-ISSN: 2349-8234  
[www.phytojournal.com](http://www.phytojournal.com)  
JPP 2022; 11(3): 62-64  
Received: 04-02-2022  
Accepted: 14-04-2022

**Precious E Oyaseiye**  
Department of Chemical  
Sciences, Niger Delta University,  
Wilberforce Island, Bayelsa,  
Nigeria

**Bamidele Martin Amos Tautua**  
Department of Chemical  
Sciences, Niger Delta University,  
Wilberforce Island, Bayelsa,  
Nigeria

**Sandile P Songca**  
School of Physics and Chemistry,  
University of KwaZulu Natal,  
Durban, South Africa

## A mini-review of pharmacological and economical values of *Funtumia africana*

**Precious E Oyaseiye, Bamidele Martin Amos Tautua and Sandile P Songca**

DOI: <https://doi.org/10.22271/phyto.2022.v11.i3a.14418>

### Abstract

*Funtumia africana* belongs to the family Apocynaceae. It is a fast growing tree found in green forests and forest reserves in West Africa. The plant has a long history of use in traditional medicine to treat and manage various diseases such as diabetes, cancer, burns, amoebic dysentery, urinary incontinence, bacterial infections and malaria. It is used economically for the production of furniture, match sticks and firewood. This review paper presents the botanical description, pharmacological uses and economic importance of *F. africana*.

**Keywords:** *Funtumia africana*, phytochemicals, medicinal plant

### Introduction

Humans have relied on plants as a source of medicine throughout history, and this is especially true in developing countries, where traditional medicine plays an important role in health care and medicinal plant research [1, 2]. Plant-based therapies have significantly improved human health and well-being [3]. The healing properties of these medicinal plants are as a result of their high content of alkaloids and polyphenolic compounds and many other bioactive phytochemicals [4]. Infusions from such plants' parts (leaves, stems, barks and roots) are used in traditional healing systems in the Niger Delta region of Nigeria in the management and treatment of diseases such as diabetes, fever, malaria, abdominal diseases, etc. [5].

### Botanical description of the plant

*Funtumia africana* (Benth) commonly known as 'Bush rubber' belongs to the family Apocynaceae, generally found in green forests and forest reserves in west Africa especially Eastern and Western part of Nigeria [6]. It is a tropical tree, usually growing up to 30 m in height (Figure. 1).

*Funtumia africana* and *Funtumia elastica* are the two species that make up the genus *Funtumia*. Both species have comparable leaves that are glabrous, leathery, elongated, elliptic more or less acuminate, cuneate at the base, and have short stalks [7]. Flowers and fruits of *Funtumia africana* are longer than those of *Funtumia elastica*. The latex of *F. elastica* coagulates easily while that of *F. africana* does not [8].

The bark of *F. africana* is brown to dark in color, thin and fissured, and then becomes granular as the tree matures. The leaves are elliptic or ovate in shape, with a round or cuneate base, acuminate apex measuring 20×9 cm, and wavy leaf margins. The primary lateral vein axels are not pitted. Flowers are yellowish white, fragrant in dense cymes with a corolla tube and lobes 6-10 mm, and 5-7 mm in length respectively. The ripe fruits are greyish brown, fusiform, with an acute or acuminate apex, up to 30 cm long with hairy wind-borne seeds [9].




Fig 1: Leaves and flowers of *F. africana* (A); *F. africana* tree (B)

**Corresponding Author:**  
**Bamidele Martin Amos-Tautua**  
Department of Chemical  
Sciences, Niger Delta University,  
Wilberforce Island, Bayelsa,  
Nigeria

### Phytochemistry of the plant

Many pharmacologically active phytochemicals are found in *Funtumia africana*, including alkaloids, flavonoids, tannins, saponin, cardiac glycosides, terpenoids, reducing sugars, steroids, and phenolic compounds [10].

Funtumine and funtumidine are the main steroidal alkaloids in *F. africana*. They are hypotensive and are used as an anesthetic locally [9]. Methyl ursolate was isolated from the chloroform fraction of the leaves of *F. africana* [11] (Figure 2).



**Fig 2:** Compounds isolated from *F. africana*

### Pharmacological uses of the plant

*Funtumia africana* is a common plant in African traditional medicine, with its many parts used to treat various diseases, such as inflammation, malaria, cancer, fever amoebic dysentery, diarrhea, and bacterial infections have all been treated with the plant, which has also been demonstrated to have larvicidal properties [12]. In West Africa, it has traditionally been used to treat urinary incontinence and burns [13].

People in Niger Delta region of Nigeria have used *F. africana* leaves for diabetes management and treatment. Findings revealed that the leaf extracts/fractions possess antioxidant and free-radical scavenging activities [14]. Antioxidants such as saponins, flavonoids, polyterpenes, polyphenols and vitamin E contains blood boosting or replenishing activities that effectively prevent excessive oxidative stress caused by certain medications and external sources. The leaf extract also inhibits lipid peroxidation and restores hematological indices to a normal level [15]. Many local traditional medicine practitioners claimed the leaves of *F. africana* have hepatoprotective activities when they were used for the treatment of hepatic diseases such as hepatitis infection [6]. The bark extract possesses broad antibiotic, antimalarial, anticancer, anti-inflammatory, and antifungal activities [11]. The root decoction is drunk for amoebic dysentery, and it can also be pounded and mixed with palm wine and water which have been used in the Benin area of Nigeria for incontinence of urine [8].

### Current Research

Over the past five years *F. africana* has been widely investigated possibly due to its phenolic content. The root extract inhibited formation of advanced glycation end-products, and lipid peroxidation [16]. These findings align with the antioxidant studies of the methanol leaf extracts which showed decrease in glutathione-peroxidase, superoxide dismutase, and catalase activities [17]. Recently, these antioxidant properties were rationalized in terms of the total

phenolic, flavonoid, and tannins contents [18]. The ethanol leaf extracts restored normal liver function in rats; regulated minerals, prostate enzymes and prostate gland morphology in rats [19, 20]. The leaf extracts also showed positive impact on rat prostatic hyperplasia indicators such as levels of testosterone, dihydrotestosterone, and estradiol and also showed bactericidal activity against strains of methicillin resistant *Staphylococcus aureus* [21, 22]. The observed moderation of triacylglycerol and high-density lipoproteins suggest that the combined *F. africana* extract possesses antilipidaemic properties [23]. Therefore, there has been escalating medicinal interest in *F. africana*.

### Economic Importance

Latex has been used as a bird lime in the past. The wood is white and even textured which can be used to make cheap joinery and furniture; it burns well and might be suitable for match sticks manufactures. The wood is also said to be utilized for crafting chairs, doors, and other household necessities [24].

The branches and trunk can be used as fuel wood. In West Africa, seed floss is used to stuff cushions. In the forest, the root system of *F. africana* protects moist soil from erosion. The littered leaves improve soil fertility in the surrounding while the fragrant yellow white flowers are visited by bees for nectar and pollen [9].

### Conclusion

*F. africana* is rich in phytochemicals, especially steroidal alkaloids and polyphenolic compounds. Extracts of this plant are utilized in the treatment and management of several diseases such as amoebic dysentery, incontinence of urine, severe burns, diabetes and malaria. *F. africana* is economically important because apart from its medicinal benefits, its branches and trunk are sources of wood used for making furniture, match sticks, and firewood.

*F. africana* has the potential of being a raw material for drug discovery which will lead to economic growth of the local communities. Further investigation of the plant is necessary to fully understand its chemical constituents and to ensure that the plant extracts are safe for consumption.

### References

1. Farnsworth NR. The role of medicinal plants in drug development. In: Krogsgaard-Larsen, S., Brogger Christensen, S., Kofod, H. (Eds), Natural Products and Drug development. Munksgaard, Copenhagen, 1994.
2. Srivastava J, Lambert J, Vietmeyer N. Medicinal plants: An Expanding Role in Development. The World Bank, Washington, DC. 1996, 18.
3. Iwu MM, Duncan AR, Okunji CO. New antimicrobial of plant origin. Janick, J (Eds), ASHS Press, Alexandria, VA, 1999.
4. Ajoko IT, Amos-Tautua BMW, Songca SP. Ethnomedicinal and economical profile of *Triumfetta cordifolia*: A mini review. Journal of medicinal plants studies. 2020;8(5):208-209.
5. Amos-Tautua BM, Kazeem AA, Olusiji A, Olatomide AF, Adebayo O, Songca SP. Effect of the leaf extracts of *Funtumia africana* (Benth.) Stapf against selected pathogens. Journal of Medicinal Plants Studies. 2020;8(4):125-129.
6. Uroko RI, Agbafor A, Nwuke CP, Uhuo EN, Nweje-Anyalowu PC, Orjiakor CA. Hepatoprotective and curative effects of methanol extract of *Funtumia africana*

- leaves against carbon tetrachloride-induced liver damage in rat. *Drug Invention Today*. 2019b;12(2):352-358.
7. Keay RWJ. Review: Trees of Nigeria.-Clarendon Press Oxford. *Nordic Journal of Botany*. 1989;11:322.
  8. Burkill HM. The useful plants of the west tropical Africa. Royal Botanic Garden. 1997;2:969.
  9. Orwa C, Mutua A, Kindt R, Jamnadass R, Anthony S. Agroforestry tree database; a tree reference and selection guide version 4.0, 2009. (<http://www.worldagroforestry.org/sites/tree-database.asp>)
  10. Gurib-Fakim A. Medicinal plants: Traditions of yesterday and drugs of tomorrow. *Molecular Aspects of Medicine*. 2006;27:1-93.
  11. Ramadwa TE, Elgorashi EE, McGawa LJ, Ahmed AS, Eloff JN. Antimicrobial, anti-inflammatory activity and cytotoxicity of *Funtumia africana* leaf extract fractions and the isolated methyl ursolate. *South African Journal of Botany*. 2017;108:126-131.
  12. Odugbemi TO, Akinsulire OR, Aibinu IE, Fabeku PO. Medicinal plants useful for malaria therapy in Okeigbo, Ondo State, Southwest Nigeria. *African Journal of Traditional & Complementary Alternative Medicine*. 2006;4:191-8.
  13. Wagner H, Seegert K, Sonnenbichler H, Ilyas M. Steroidal alkaloids of *Funtumia africana*. *Plant Medica*. 1987;53(5):444-9.
  14. Amos-Tautua B, Ajileye O, Ndoni S, Bamidele F, Onigbinde A. Evaluation of phenolic contents, free radical scavenging activity and functional group analysis of the leaf extract of a medicinal plant in Niger Delta Region. *Chemistry International*. 2017;3:250.
  15. Uroko RI, Egba SI, Uchenna ON, Orjiakor CA, Agbafor A, Alaribe CC. Therapeutic effects of methanolic extract of *Funtumia africana* leaves on antioxidants and haematological indices of carbon tetrachloride-induced oxidative stress in rats. *Drug Invent Today*. 2019a;12(1):114-122.
  16. Nkemehule FE, Ogonnia SO, Odukoya OA. Antiglycation and antioxidant activity of *Funtumia africana* root used in the management of diabetes. *Tropical Journal of Natural Product Research*. 2018;2(5):203-208. <https://doi.org/10.26538/tjnpr/v2i5.1>
  17. Ikechukwu UR, Egba SI, Uchenna ON, Orjiakor CA, Agbafor A, Alaribe CC. Therapeutic effects of methanolic extract of *Funtumia africana* leaves on antioxidants and hematological indices of carbon tetrachloride-induced oxidative stress in rats. *Drug Invention Today*. 2019;12(1):114-122.
  18. Frempong TF, Badu M, Boamah VE, Amponsah IK, Agbemade B, Boateng RA, *et al*. Phenolic content, antioxidant properties and antimicrobial activities of the extracts from *Funtumia africana* and *Funtumia elastica*. *Chemistry Africa*. 2021; 4: 503–512. <https://doi.org/10.1007/s42250-021-00249-5>
  19. Uroko RI, Adamude FA, Egba SI, Chukwu CN, Asadu CL, Okwara EC. Effects of combined ethanol extract of *Funtumia africana* and *Abutilon mauritianum* leaves (FAAM) on liver function indices of benign prostatic hyperplasia (BPH) induced rats. *Herba Polonica*. 2020a;66(3):24-35. <https://doi.org/10.2478/hepo-2020-0013>.
  20. Uroko RI, Adamude FA, Egba SI, Nwuke CP, Asadu CL, Anyaorah P. Effects of combined ethanol extract of *Funtumia africana* and *Abutilon mauritianum* leaves on prostate biomarkers and serum mineral levels in prostatic hyperplasia induced male rats. *Journal of Renal Endocrinology*. 2020b;7(1):e06-e06. <https://doi.org/10.34172/jre.2021.06>
  21. Uroko RI, Nwuke CP, Egba SI, Uchenna ON, Nwaekwu FE, Anyanwu PO. Effects of combined ethanolic extracts of *Funtumia africana* and *Abutilon mauritianum* leaves on sex hormones and prostate indices of benign prostatic hyperplasia induced rats. *Animal Research International*. 2020c;17(3):3892-3902.
  22. Kouadio NJ, Kipre BG, Guessenn NK, Kouassi KA, Yao K, Kone MW. Activity evaluation of leaves' extracts of *Funtumia africana* Stapf (Apocynaceae) towards Methicillin-Resistant *Staphylococcus aureus* (MRSA) and phytochemical screening. *SM Journal of Medicinal Plant Studies*. 2017;1(1):1001. <https://doi.org/10.36876/smjmps.1001>
  23. Uroko RI, Chukwu CN, Egba SI, Adamude FA, Ajuzie JC. Combined ethanol extract of *Funtumia africana* and *Abutilon mauritianum* leaves improves the lipid profile and kidney function indices of benign prostatic hyperplasia in rats. *Acta Scientiarum Polonorum, Technologia Alimentaria*. 2020d;19(4):395-404. <https://doi.org/10.17306/J.AFS.0819>
  24. Palgrave KC. *Trees of Southern Africa*. 3rd edition, New Holland Publishing, South Africa, 2002, 958.